

Journal of Plant Protection and Pathology


Journal homepage & Available online at: www.jppp.journals.ekb.eg

Identification of the Resistance Gene *Sr2* in some Egyptian Wheat Hybrids

Mona E. Mohamed^{1*} and Shaimaa E. Ibrahim²

¹ Plant Pathology Research Institute, Wheat Diseases Research Dept., A.R.C. Egypt, 12619.

² Wheat Res. Dept., Field Crops Res. Institute, ARC, Egypt, 12619.



ABSTRACT

Seven Egyptian wheat cultivars were crossed with *Sr2* (the stem rust resistant gene), and the parents, F₁ and F₂ crosses were tested to natural stem rust disease in the field of Gemmeiza Agricultural Station, ARC, during the 2019 to 2021 growing seasons. The *Sr2* gene was completely resistant to stem rust, while the Egyptian cultivars showed different responses to the disease. Sakha-95 was completely resistant, Misr-1 and Misr-2 showed high susceptibility. Gemmeiza-12, Giza-171, and Sakha-94 showed moderate resistance (MR), while a trace of moderate susceptibility (Tr-MS) was recorded for the cultivar Misr-3. All the F₁ crosses were resistant, indicating the dominance of resistance to stem rust disease. Chi-square goodness of fit was used to assess the independence of *Sr2* resistance gene in the tested F₂ crosses. The crosses *Sr2*XGemmeiza-12, *Sr2*X Sakha-94, *Sr2*X Sakha-95 and *Sr2*X Giza-171 have the expected segregation ratio of 15:1 for a duplicate gene. However, the crosses *Sr2*X Misr-1 and *Sr2*X Misr-3 fit a 3:1 ratio conditioned by one dominant gene. The resistance of the cross *Sr2*X Misr-2 fits complementary gene action (9:7). Three specific SSR markers (*xgwm533*, *stm559tgag*, and *stm598cac*) were used to identify *Sr2* in the Egyptian cultivars and their F₁ hybrids. The three markers confirmed presence of *Sr2* gene in the Egyptian cultivars and their F₁ hybrids. However, the *stm598cac* marker could differentiate between the tested cultivars and their F₁ hybrids by amplifying an additional band with a molecular weight of 200bp. Molecular analysis confirmed the results of chi-square test and could be serving in developing wheat programs.

Keywords: wheat, stem rust, segregating population

INTRODUCTION

Stem rust caused by *Puccinia graminis* f. sp. *tritici* is among the fungal diseases that significantly affect wheat production around the world (Bhavani *et al.*, 2022; Guo *et al.*, 2022). To combat stem rust, breeding for resistant cultivars remains the most efficient and reliable method to control this disease (Kumar *et al.*, 2022b). There are different sources of resistance to stem rust, such as wheat wild relatives. *Triticum turgidum* (Yaroslav emmer) was transferred to the Hope cultivar (Mago *et al.*, 2011). Slow rusting genes like *Sr2*, *Sr55*, *Sr56*, *Sr57*, and *Sr58* confer resistance at the adult plant stage under high disease pressure (Herrera-Foessel *et al.*, 2014; Singh *et al.*, 2015; Kosgey *et al.*, 2021). Previously, *Sr2* was used as a parent in several breeding programs around the world and used frequently in CIMMYT breeding materials (Singh *et al.*, 2011a). There are several sources for *Sr2* gene, including Pavon, Arthur, Lancer, and Hope. When present alone, the *Sr2* gene showed slow rusting under intense disease pressure, but when combined with other minor genes to form the "Sr2 complex," it provides an appropriate level of resistance (Singh *et al.*, 2011a; Singh *et al.*, 2015). This gene tends to be non-specific and is currently effective against all isolates of *Puccinia graminis* f. sp. *tritici* throughout the global regions that grow wheat. In Egypt, *Sr2* (Pavon76 cultivar) has shown stable adult-plant resistance (APR) to stem rust in the field from 2016–2020 seasons (El-Shamy *et al.*, 2019; Elkot *et al.*, 2020). Shahin *et al.* (2020) evaluated *Sr2*, *Sr24*, and *Sr26* under field conditions during the 2014 and 2015 seasons in Kafr elsheikh, Sharqia, and Nubaria governorates and found that *Sr2* was highly effective for

resistance in both seasons. El-Shamy *et al.* (2019) found that Pavon showed complete resistance to stem rust under field conditions at recommended and late dates of cultivation. Molecular genetic analysis has been used for the evaluation of both major and minor genes in various crops. Moreover, in wheat, molecular markers linked to race-specific genes for rust resistance have been identified and utilized for marker-assisted selection (Suenaga *et al.*, 2003; Javadi *et al.*, 2021). Molecular markers have been developed for numerous stem rust resistance genes, including *Sr2* (Spielmeyer *et al.*, 2003; Hayden *et al.*, 2004; Mago *et al.*, 2011). The aims of this study are to detect gene actions of resistance against stem rust disease in seven crosses with *Sr2* gene and to detect the presence of *Sr2* gene in the seven Egyptian bread wheat cultivars using molecular marker technique.

MATERIALS AND METHODS

1. Plant materials

Seven of the recent Egyptian bread wheat cultivars i.e. Gemmeiza-12, Sakha-94, Sakha-95, Giza-171, Misr-2, Misr-3 and Misr-1 were crossed as females with the resistant cultivar Pavon76 carrying stem rust gene *Sr2* as a male parent in 2018/ 2019 growing season at Gemmeiza Agricultural Station, ARC. The names and pedigree of tested materials were reported by El-Shamy *et al.* (2019). Seeds of Pavon (*Sr2*) stock were obtained from ICARDA, while the bread wheat cultivars derived from The National Wheat Dept., Crop Research Institute. In 2019/2020 growing season, the hybrid seeds were sown in 1row, 2m long, 40 cm apart and 20 cm within rows for each cross to have F₁ seeds. In 2020/2021 growing season, the parents and F₁ hybrid seeds were sown in

* Corresponding author.

E-mail address: mmonaelsayed@yahoo.com

DOI: 10.21608/jppp.2022.168076.1106

1row, 2m long, 40 cm apart and 20 cm within rows while about 100 to 150 grains of F2 seeds of each cross were drilled in plots consisted of 6 rows, 7m. long, 20cm apart and 10 cm in-between seeds for evaluation to stem rust disease. The experiment was surrounded by border 1 m of a highly susceptible cultivar i.e. Morocco and left to natural stem rust infection. Fertilizer application followed recommended cultural practices 70 kg N and 100 kg P/fed.

2. Disease assessment

Disease severity was scored for the parents, F1 and F2 plants of each cross when Morocco plants showed maximum rust severity using the modified Cobb scale (Peterson *et al.*, 1948) and the host response was classified as either 0 (immune), R (resistant), MR (moderately resistant), M (intermediate), MS (moderately susceptible) or S (susceptible) according to Roelfs (1992). The disease severity is multiplied by a constant value for each infection type of stem rust to calculate (ACI) according to (Saari and Wilcoxon, 1974), where: R= 0.2, MR = 0.4, MS = 0.8, S = 1.00.

3. Chi-square analysis

Chi-square analysis was used to evaluate the goodness of fit of observed segregation with expected ratios and to

Table 1. Sr gene, Markers, expected PCR fragment size, and annealing temperature.

Sr gene	Markers	Expected fragment size (bp)	Primer sequence	Annealing temperature	References
Sr 2	xgwm533	120bp	GTTGCTTACGGGAAAGCC AAGGCAGAATCAAACCGGAATA	48.3°C	Hayden <i>et al.</i> , 2004
	stm559tgag	from 79 to 85 bp	AAGGCAGAATCAAACCGGAATA TGTGTGTGTGTGAGAGAGAG	50°C	
	stm598cac	from 61 to 67 bp	GTTGCTTACGGGAAAGCC TCTCTCTCTC TCTCACACACAC	49.3°C	

RESULTS AND DISCUSSION

Results

Evaluation at the adult stage

The *Sr2* showed completely resistance to stem rust under field conditions at the adult stage during the years of the study (Fig.1). The Egyptian wheat cultivars showed varied reactions to stem rust disease under field conditions ranging from zero to susceptible reaction (30S). The wheat cultivars divided into two groups, the resistant group was Sakha-95(0), Gemmeiza-12(Tr- MR), Giza-171(Tr-MR), and Sakha-94 (5MR). While the susceptible group included Misr-3 (Tr-MS), Misr-2(20S), and Misr-1(30-S), respectively.




Fig. 1. Mean disease severity of 7 Egyptian bread wheat cultivars and *Sr2* to stem rust disease during the growing season 2020/2021.

The Chi-square test

An analysis of adult plants resistance of F1 and F2 populations to stem rust disease is shown in Table (2). The results clear that all the F1 hybrids of the seven crosses showed adult plant resistance responses to natural infection of

estimate the number of genes responsible for stem rust resistance of the F2 population (Snedecor and Cochran 1989).

4. Molecular study

DNA extraction

The total DNA of *Sr2*, wheat cultivars, and their F1 hybrids were extracted from 200 mg of 10 days seedlings. The extraction method was carried out according to Emara *et al.* (2016).

PCR amplification conditions

Three specific microsatellites (SSR) markers linked to *Sr2* were used to detect its presence in the 7 bread wheat cultivars and their F1 hybrids. The markers' names, sequences, and annealing temperatures were listed in Table 1. The PCR was conducted following Hayden *et al.* (2004). Each PCR mixture has the following composition; Promega Nuclease free water (9.5 μ l), 25ng nucleic acid (1 μ l), each primer (10) pmol (1 μ l), and GoTag® Colorless Master Mix 12.5 μ l, PCR products (25 μ l) were analyzed by electrophoresis in a 2% agarose gel, stained with ethidium bromide (7.0 μ g/50 ml) and DNA bands were visualized using a UV trans-illuminator and photographed.

stem rust. The F2 populations showed segregation in its responses as follow. The crosses *Sr2*XGemmeiza-12, *Sr2*X Sakha-94, *Sr2*XSakha-95 and *Sr2*XGiza-171 showed 15R:1S ratio, which segregated in 150R:10S, 220R:20S, 200R:30S, and 230R:10S, respectively. However, F2 plants of the crosses *Sr2*XMisr-1 and *Sr2*XMisr-3 showed 3:1 ratio which segregated in 110R:30S and 160R:50S, respectively. The F2 plants of the cross *Sr2*XMisr-2 showed 9R:7S which segregated in 80R:60S.

The molecular detection of *Sr2*

The markers xgwm533, stm559tgag, and stm598cac were used to determine the presence of *Sr2* in the Egyptian cultivars and the F1 hybrids. Amplification with the xgwm533 primer developed for *Sr2* produced two PCR amplicon with a size of 120 bp and 65 bp (Fig. 2). The amplified fragment with 120 bp was present in all of the tested cultivars and the F1 hybrids. While, the additional band with molecular weight from 65 bp appeared in all the Egyptian cultivars, and absent from their F1. Amplification with the stm559tgag marker developed for *Sr2* produced a single PCR amplicon with a size about 80 bp in the *Sr2* parent but in the Egyptian cultivars and their F1 hybrids with the molecular size 100 p (Fig. 3). Amplification with the stm598cac primer produced two amplicons with size from 61 to 67 bp in the *Sr2*, the Egyptian cultivars as parents and their F1 hybrids (Fig. 4). Another fragment was amplified with molecular weight 200bp in Misr-1, Misr-3 and Giza-171 and their F1 hybrids. Moreover, this band appears in Sakha-95 cultivar and absent from its F1 hybrids. However, it was absent in Gemmeiza-12, Sakha -94, and Misr-2 cultivars and present in their F1 hybrids (Table 3).

Table 2. Segregation of resistance to stem rust pathogen in the F2 populations derived from the cross Sr2 and 7 Egyptian wheat cultivars.

Cross name	No. of tested plants	Response			Segregation		Expected ratio		χ^2	p-value
		*R	**I	***S	R	S	R	S		
Sr2XGemmeiza 12	F ₁	54	38	16						
	F ₂	160	130	20	10	150	10	15	1	0.00
Sr2X Sakha-94	F ₁	65	44	21						
	F ₂	240	200	20	20	220	20	15	1	1.778
Sr2XSakha- 95	F ₁	49	40	9						
	F ₂	230	190	10	30	200	30	15	1	18.116
Sr2XMisr-1	F ₁	52	33	19						
	F ₂	140	80	30	30	110	30	3	1	0.952
Sr2XMisr-2	F ₁	44	30	14						
	F ₂	140	40	40	60	80	60	9	7	0.045
Sr2XMisr-3	F ₁	47	34	13						
	F ₂	210	110	50	50	160	50	3	1	0.159
Sr2XGiza-171	F ₁	50	39	11						
	F ₂	240	130	100	10	230	10	15	1	1.778

*R= Resistant, **I= Intermediate, ***S= Susceptible




Fig. 2. xgwm533-PCR amplifications in Sr2 (120 bp), 7 wheat cultivars, and their F1 hybrids. 1: 50bp DNA ladder RTU (Gene Direx), 2: Sr2 3: Gemmeiza-12, 4: Sr2 x Gemmeiza-12, 5: Sakha-94, 6: Sr2 XSakha-94, 7: Sakha-95, 8: Sr2 X Sakha-95, 9: Misr-1, 10: Sr2 X Misr-1, 11: Misr-2, 12: Sr2 XMisr-2, 13: Misr-3, 14: Sr2 X Misr-3, 15: Giza-171, 16: Sr2X Giza-171, 17: Sr2.




Fig. 3. stm559ttag PCR amplifications in Sr2 (100 bp), 7 Egyptian wheat cultivars and their F1 hybrids. 1: 50bp DNA ladder RTU (Gene Direx), 2: Sr2 3: Gemmeiza-12, 4: Sr2 x Gemmeiza-12, 5: Sakha-94, 6: Sr2 XSakha-94, 7: Sakha-95, 8: Sr2 X Sakha-95, 9: Misr-1, 10: Sr2 X Misr-1, 11: Misr-2, 12: Sr2 XMisr-2, 13: Misr-3, 14: Sr2 X Misr-3, 15: Giza-171, 16: Sr2X Giza-171, 17: Sr2.




Fig. 4. stm598cac-PCR amplifications in Sr2 (from 61 to 67 bp), 7 Egyptian wheat cultivars, and their F1 hybrids. 1: 50bp DNA ladder, 2: Sr2 3: Gemmeiza-12, 4: Sr2 x Gemmeiza-12, 5: Sakha-94, 6: Sr2 XSakha-94, 7: Sakha-95, 8: Sr2 X Sakha-95, 9: Misr-1, 10: Sr2 X Misr-1, 11: Misr-2, 12: Sr2 XMisr-2, 13: Misr-3, 14: Sr2 X Misr-3, 15: Giza-171, 16: Sr2X Giza-171, 17: Sr2.

Table 3. Presence/absence of Sr2 in Egyptian cultivars and their F1 hybrids.

Cultivars	xgwm533		stm559ttag		stm598cac	
	120bp	65bp	100 bp	61-67bp	200bp	
Gemmeiza-12	1	1	1	1	1	0
Sr2XGemmeiza-12	1	0	1	1	1	1
Giza-171	1	1	1	1	1	1
Sr2X Giza-171	1	0	1	1	1	1
Sakha-94	1	1	1	1	0	0
Sr2X Sakha-94	1	0	1	1	1	1
Sakha-95	1	1	1	1	1	1
Sr2X Sakha-95	1	0	1	1	0	0
Misr-1	1	1	1	1	1	1
Sr2X Misr-1	1	0	1	1	1	1
Misr-2	1	1	1	1	0	0
Sr2XMisr-2	1	0	1	1	1	1
Misr-3	1	1	1	1	1	1
Sr2X Misr-3	1	0	1	1	1	1

1=present 0=absent

Discussion

Wheat stem rust affects wheat production around the world as more than 90% of wheat cultivars have susceptible reactions to the Ug99 race group (Singh *et al.*, 2011b). In Egypt, this pathogen causes 1.96 to 8.21% losses in the yield of most of the local wheat varieties (Shehab-Eldeen and Abou-Zeid, 2020). The deployment of diverse resistance genes especially from wild relatives is the most effective way to reduce the potential of stem rust. To date, 73 *Sr* genes/alleles have been identified and about 35 are effective against the Ug99 race lineages (Singh *et al.*, 2015). Unfortunately, most of them are race specific and don't provide durable resistance to stem rust. On the other hand, adult plant resistance provides long-lasting durable resistance to multipathogenic resistance and is provided by some resistant genes including *Sr2*, *Sr54*, *Sr55*, and *Sr57* (Kosgey *et al.*, 2021). Of them, *Sr2* has been successfully deployed in many breeding programs around the world like Australia, India, CIMMYT, and China due to its high yield and resistance to stem rust (McIntosh 1995; Malik *et al.*, 2013, Bhavani *et al.*, 2014; Xu *et al.*, 2017). In Egypt, *Sr2* has been evaluated under field condition and provide high resistance to stem rust (El shamy *et al.*, 2019; Shahin *et al.*, 2020; Elkot *et al.*, 2016; El-Orabey *et al.*, 2019). Also, Singh *et al.* (2006) reported that both Sonalika and Pavon 76 (*Sr2*) were resistant during field assessments in 2004 and 2005 in Kenya with a maximum disease score of 15MS. In our study *Sr2* was crossed as a male parent with seven Egyptian cultivars, and their F1 and F2 were evaluated to stem rust under natural infection during the study. *Sr2* and Sakha-95 were completely resistant to stem rust. Gemmeiza-12, Giza-171 and Sakha-94 showed moderate resistance ranged from traces to 5 MR. While cultivars Misr-3,

Misr-2, and Misr-1 showed susceptible reaction ranged from traces MS to 30-S. The F1 of all crosses showed completely resistance to stem rust indicating that the resistance is dominant, but the F2 population showed segregation in their responses to the disease. A chi-square goodness-of-fit test was calculated for F2 crosses to determine the number of genes that control resistance to stem rust. The chi-square test on the observed segregation ratio was consistent with digenic Mendelian segregation for the crosses *Sr2XGemmeiza-12*, *Sr2XSakha-94*, *Sr2XSakha-95* and *Sr2XGiza-171* segregated in 15R:1S ratio, indicating that two dominant independent genes conditioning the resistance to stem rust in these crosses. However, the F2 plants of the crosses *Sr2XMisr-1* and *Sr2XMisr-3* segregated in 3:1 ratio indicating that the two crosses possess one (a single) dominant gene. The F2 plants of the cross *Sr2XMisr-2* showed 9R:7S which segregated in 160R:50S indicating that it possess a pair of dominant complementary genes. Nzuve *et al.* (2013) crossed five resistant wheat lines (KSL-2, 3, 5, 12 and 19) with the susceptible line Cacuke to determine the stem rust resistance gene inheritance. Chi square test showed that the segregation in KSL-2 parent is conditioned by a single dominant gene of resistance. While, the stem rust inheritance in the parents KSL-5, KSL-19, KSL-12 and KSL-3 were conditioned by two genes. Previous studies conducted in Egypt showed that *Sr2* provides resistance to stem rust under field conditions and the tested Egyptian cultivars contain *Sr2* as confirmed by SSR markers (Elkot *et al.*, 2020; Abu Aly *et al.*, 2014). However, it couldn't differentiate between cultivars regardless of their response to stem rust-resistant and susceptible ones. In our study, three diagnostic-specific markers linked to *Sr2* were used; gwm533, stm559ttag, and stm598cac to accurately determine the presence of the *Sr2* gene in the Egyptian cultivars and their F1 hybrids. The two additional markers stm559ttag and stm598cac were used in this study to overcome the drawbacks of the gwm533 marker. Hayden *et al.* (2004) stated that *Sr2* gene closely linked microsatellite marker Xgwm533 and typically amplifies a 120-bp fragment from wheat lines known to carry *Sr2*. However, it couldn't discriminate between the *Sr2* carrier and non-*Sr2* carrier wheat lines. Also, Vishwakarma *et al.* (2019) suggested the existence of allelic homoplasy in *Sr2* gene non-carriers varieties like the Aroona variety. The three tested markers indicated the presence of *Sr2* in the tested cultivars and their crosses. The stm598cac marker is considered a diagnostic marker as it could differentiate between tested cultivars and their F1 hybrids by amplifying an additional band with a molecular weight of 200bp. This maker is considered as a diagnostic marker since it could make differences between the tested cultivars.

ACKNOWLEDGMENT

The molecular study was carried out at EPCRS Excellence Center certified Plant Pathology and Biotechnology Lab, Department of Agricultural Botany, Faculty of Agriculture, Kafr ElSheikh University, Egypt.

CONCLUSION

Sr2 is a good source of resistance to stem rust since it confers resistance to stem rust. Moreover, the stm598cac marker is a diagnostic marker as it is the only marker that could differentiate between the tested cultivars and their F1 hybrids by amplifying an additional band with a molecular weight of 200bp.

REFERENCES

- Abu Aly, A. A. M., Shahin A.A., EL-Naggar, D. R. and Hermas, G. 2014. Identification and evaluation of resistance genes sources of stem rust in different Egyptian and CIMMYT Wheat genotypes using conventional and molecular techniques. *J. Plant Prot. and Path.*, Mansoura Univ., 5 (6): 729 – 70.
- Bhavani, S., Hodson, D.P., Huerta-Espino, J., Randhawa, M.S. and Singh, R.P. 2019. Progress in breeding for resistance to Ug99 and other races of the stem rust fungus in CIMMYT wheat germplasm. *Front. Agr. Sci. Eng.*, 6(3): 210–224.
- Bhavani, S., Singh, R.P., Hodson, D.P., Huerta-Espino, J. and Randhawa, M.S. 2022. Wheat Rusts: current status, prospects of genetic control and integrated approaches to enhance resistance durability. In *Wheat Improvement* (pp. 125-141). Springer, Cham.
- Elkot, A.F.A., Abd El-Aziz, M.H., Aldrussi, I.A. and El-Maghrary, M.A. 2016. Molecular identification of some stem rust and yellow rust resistance genes in Egyptian wheat and some exotic genotypes. *Ass. J. Agric. Sci.*, 47:124-135.
- Elkot, A. F., El-Orabey, W. M., Draz, I. S. and Sabry, S. R. 2020. Marker-assisted identification of stem rust resistance genes *Sr2*, *Sr13*, *Sr22* and *Sr24* in Egyptian wheat cultivars. *Egypt. J. Plant Breeding*, 24(1): 225-245.
- El-Orabey, W.M., Elbasyoni I.S., El-Moghazy S.M., Ashmawy, M.A. 2019. Effective and ineffective of some resistance genes to wheat leaf, stem and yellow rust diseases in Egypt. *J. Plant Production*, Mansoura Univ., 10 (4): 361-371.40, pp.279-287.
- El-Shamy, M.M., Elmassry, E.L. and Sallam, M. E.A. 2019. Line x tester mating design analysis for grain yield, yield attributes and stem rust disease resistance in bread wheat. *Egypt. J. Plant Breeding*, 23(5): 971 – 993.
- Emara, H.M., Omar, A.F., El-Shamy, M.M. and Mohamed, M.E. 2016. Identification of *Pm24*, *Pm35* and *Pm37* in thirteen Egyptian bread wheat cultivars using SSR markers. *Ciência e Agrotecnologia*, 40:279-287.
- Guo, Y., Betzen, B., Salcedo, A., He, F., Bowden, R.L., Fellers, J.P., Jordan, K.W., Akhunova, A., Rouse, M.N., Szabo, L.J. and Akhunov, E. 2022. Population genomics of *Puccinia graminis* f. sp. *tritici* highlights the role of admixture in the origin of virulent wheat rust races. *Nature communications*, 13(1):1-16.
- Hayden, M.J., Kuchel, H. and Chalmers, K.J. 2004. Sequence tagged microsatellites for the Xgwm533 locus provide new diagnostic markers to select for the presence of stem rust resistance gene *Sr2* in bread wheat (*Triticum aestivum* L.). *Theoretical and Applied Genetics*, 109(8):1641-1647.
- Herrera-Foessel, S.A., Singh, R.P., Lillemo, M., Huerta-Espino, J., Bhavani, S., Singh, S., Lan, C., Calvo-Salazar, V. and Lagudah, E.S. 2014. Lr67/Yr46 confers adult plant resistance to stem rust and powdery mildew in wheat. *Theoretical and Applied Genetics*, 127(4):781-789.
- Javadi, H., Dadkhodaie, A. and Heidari, B., 2021. Molecular marker analysis of stem rust resistance genes in some Iranian wheat genotypes. *Cytology and Genetics*, 55(5):460-470.

- Kosgey, Z.C., Edae, E.A., Dill-Macky, R., Jin, Y., Bulbula, W.D., Gemechu, A., Macharia, G., Bhavani, S., Randhawa, M.S. and Rouse, M.N. 2021. Mapping and validation of stem rust resistance loci in spring wheat line CI 14275. *Frontiers in plant science*, 11, p.609659.
- Kumar, A., Choudhary, A., Kaur, H. and Mehta, S., 2022b. A walk towards Wild grasses to unlock the clandestine of gene pools for wheat improvement: A review. *Plant Stress*, 3, p.100048.
- Mago, R., Brown-Guedira, G., Dreisigacker, S., Breen, J., Jin, Y., Singh, R., Appels, R., Lagudah, E.S., Ellis, J. and Spielmeyer, W. 2011. An accurate DNA marker assay for stem rust resistance gene *Sr2* in wheat. *Theoretical and applied genetics*, 122(4):735-744.
- Malik, R., Parveen, S., Saharan, M.S., Kumar, R., Sharma, A.K., Bhardwaj, S.C. and Sharma, I. 2013. Characterization of stem rust resistance gene *Sr2* in Indian wheat varieties using polymerase chain reaction (PCR) based molecular markers. *African Journal of Biotechnology*, 12(18).
- McIntosh, R.A., Wellings, C.R. and Park, R.F. 1995. *Wheat rusts: an atlas of resistance genes*. CSIRO publishing.
- Nzuve, F. M., Tusiime, G., Bhavani, S., Njau, P. N. and Wanyera, R. 2013. Studies of the genetics of inheritance of stem rust resistance in bread wheat. *Afr. J. Biotechnol.*, 12(21): 3133 – 3159.
- Peterson, R.F., Campbell, A.B. and Hannah, A.E. 1948. A diagrammatic scale for estimating rust intensity on leaves and stems of cereals. *Canadian journal of research*, 26(5): 496-500.
- Roelfs, A.P. 1992. *Rust diseases of wheat: concepts and methods of disease management*. Cimmyt.
- Saari, E.E. and Wilcoxson, R.D. 1974. Plant disease situation of high yielding durum wheat in Asia and Africa. *Arr. Rev. Phytopatholgy*, 2: 47- 68.
- Shahin, A., Youssif, W. and EL-Naggar, D. 2020. Detection of Ug99 (TTKSK) of wheat stem rust fungus and new virulence races of *Puccinia graminis* f. sp. *tritici* in Egypt. *Egyptian Journal of Phytopathology*, 48(1):14-28.
- Shehab-Eldeen, M.T. and Abou-Zeid, M.A. 2020. Quantitative studies on resistance to stripe and stem rust diseases and on grain yield of bread wheat. *Journal of Plant Production*, 11(11):1071-1075.
- Singh, R.P., Hodson, D.P., Huerta-Espino, J., Jin, Y., Bhavani, S., Njau, P., Herrera-Foessel, S., Singh, P.K., Singh, S. and Govindan, V. 2011a. The emergence of Ug99 races of the stem rust fungus is a threat to world wheat production. *Annu. Rev. Phytopathology*, 49:465-481.
- Singh, R. P., Hodson, D. P., Jin, Y., Huerta-Espino, J., Kinyua, M., Wanyera, R., Njau, P., and Ward, R. W. 2006. Current status, likely migration and strategies to mitigate the threat to wheat production from race Ug99 (TTKS) of stem rust pathogen. *CAB Rev. Perspect. Agric. Vet. Sci. Nutr. Nat. Resour.* 1:1-13.
- Singh, R.P., Hodson, D.P., Jin, Y., Lagudah, E.S., Ayliffe, M.A., Bhavani, S., Rouse, M.N., Pretorius, Z.A., Szabo, L.J., Huerta-Espino, J. and Basnet, B.R. 2015. Emergence and spread of new races of wheat stem rust fungus: continued threat to food security and prospects of genetic control. *Phytopathology*, 105(7):872-884.
- Singh, R.P., Huerta-Espino, J., Bhavani, S., Herrera-Foessel, S.A., Singh, D., Singh, P.K., Velu, G., Mason, R.E., Jin, Y., Njau, P. and Crossa, J. 2011b. Race non-specific resistance to rust diseases in CIMMYT spring wheats. *Euphytica*, 179(1):175-186.
- Snedecor, G.W. and Cochran, W.G. 1989. *Statistical Methods* (8th Ed.). Ame: Iowa Stae Univ. Press.
- Spielmeyer, W., Sharp, P.J. and Lagudah, E.S. 2003. Identification and validation of markers linked to broad-spectrum stem rust resistance gene *Sr2* in wheat (*Triticum aestivum* L.). *Crop Science*, 43(1):333-336.
- Suenaga, K., Singh, R.P., Huerta-Espino, J. and William, H.M. 2003. Microsatellite markers for genes *Lr34/Yr18* and other quantitative trait loci for leaf rust and stripe rust resistance in bread wheat. *Phytopathology*, 93:881–890.
- Vishwakarma, G., Sanyal, R. P., Shitre, A., Gadekar, D. A., Saini, A., and Das, B. K. (2019). Validation and marker-assisted selection of stem rust resistance gene *Sr2* in Indian wheat using gel-based and gel-free methods. *Journal of Crop Science and Biotechnology*, 22(4): 309-315.
- Xu, X.; Li, D.; Liu, Y.; Gao, Y.; Wang, Z.; Ma, Y.; Yang, S.; Cao, Y.; Xuan, Y.; Li, T. 2017. Evaluation and identification of stem rust resistance genes *Sr2*, *Sr24*, *Sr25*, *Sr26*, *Sr31* and *Sr38* in wheat lines from Gansu Province in China. *PeerJ*, 5, e4146.

تعريف جين Sr2 المقاوم لمرض الصدا الأسود في بعض هجن القمح المصريية

من السيد محمد¹ و شيماء الدسوقي ابراهيم²

¹ قسم بحوث أمراض القمح - معهد بحوث أمراض النباتات - مركز البحوث الزراعية - مصر
² قسم بحوث القمح - معهد بحوث المحاصيل الحقلية - مركز البحوث الزراعية - مصر

الملخص

تم تهجين سعة أصناف من القمح المصري مع سلاله قمح أحادية الجين (*Sr2*) تحت ظل الظروف الحرارية وتقييم الأناء والجيلين الأول والثاني لمرض صدا الساق في محطة بحوث الجميرة خلال مواسم النمو 2019 و 2021، وتم اختبار هذه النباتات في مرحلة الطور البالغ تحت ظروف العدوى الطبيعية في الحال. أثناء الدراسة كان *Sr2* مقاوماً تماماً لصدأ الساق، بينما أظهرت الأصناف المصرية رجات مقاولته من ردود الفعل لمرض الصدا الأسود. حيث كان صنف سخا-95 مقاولاً تماماً بينما الأصناف مصر-1 و مصر-2 كانت أكثر الاصناف إصابة بينما كانت الأصناف جيزة-171 و جيزة-12 و سخا-94 متوسطة المقاومة للصدأ الأسود ومصر-3 قابل لللاصبة (Tr-MS). أظهر الجيل الاول مقاومة للصدأ الساق بينما كان هناك انعزالت في هجن نباتات الجيل الثاني. استخدم اختبار مربع كائي للتعرف على عدد الجينات التي تحكم صفة المقاومة لمرض. أظهرت النتائج وجود اثنين من الجينات المستقلة تحكم صفة المقاومة في الهجن الخالصة بالجين *Sr2* مع كل من جيزة-12 او سخا-94 و سخا-95 و جيزة-171. بينما يوجد جين واحد سائد يتحكم صفة المقاومة الخالصة بالجين *Sr2* مع الأصناف مصر-1 و مصر-3. كما يوجد زوج من الجينات المكملة المهيمنة على صفة المقاومة في الهجين الخالص بالجين *Sr2* مع الصنف مصر-2. تم استخدام ثلاثة من المعلومات SSR وهي (stm598cac و stm559tgag و xgwm533) لاختبار وجود الجين *Sr2* في الأصناف المصرية وكذا في هجن الجيل الأول. كما أظهر المعلم *stm598cac* كفاءة متخصصة في التفريق بين الأصناف المختلفة وهجن الجيل الأول بوجود band متخصصة يوزن جزيئي (200bp).

الكلمات الافتتحية: القمح ، صدا الساق ، اجيال العزالية