

Effect of Magnetic Field on Seed Viability and Insect Infestation of Some Wheat Varieties

Hozayn, M.¹; Doaa M. Boraei² and Amal A. A. EL-Mahdy³

¹ Field Crop Research Department, Agricultural and Biological Research Division, National Research Centre, Giza, Egypt.

² Stored Products and Grains Department, Plant Protection Research Institute, Agriculture Research Center, Dokki, Egypt.

³ Seed Technology Research Department, Field Crops Research Institute, Agricultural Research Centre, Giza, Egypt



ABSTRACT

The effect of magnetic field periods (1 min, 6 and 12 h) exposure of some infected wheat seed varieties (Masr-1, Sids-12 and Sakha-93) with two stored grain insects *Sitophilus oryzae* (L.) and *Rizopertha dominica* F. to low static magnetic field (30&60 mT) on mortality (%), reduction in F₁ progeny (%), seeds germination (G%), seed and seedling vigor traits were studied under controlled laboratory conditions. Results showed that, the mortality percentage of both insects increased with increasing of MFs levels and time exposure. Magnetic field level at 60 mT for 12 hrs was the most effective against tested insects in which mortality percentage of *S. oryzae* was 56.60, 53.30 and 50.00 % with Masr-1, Sakha-93 and Sids-12 respectively after 10 days from exposure period while at the same level and time, mortality percentage of *R. dominica* was 45.00, 36.60 and 33.30% with Sids-12, Sakha-93 and Masr-1, respectively. Exposure infected seeds with (*S.oryzae* and *R.dominica*) to magnetic field treatments 60 mT for (1 min.,6 and 12 h) positively affected on all seed and seedling vigor parameters compared to control treatment (infected seed without magnetic treatment). Moreover almost equaled with uninfected seeds but it showed significant effect on all parameters comparing with untreated seeds. The results suggest that pre-sowing low-magnetic field treatment has the potential to improve seed germination and seedling vigor traits of infected wheat varieties with stored grain insects(*S. oryzae* or *R. dominica*) through reduction in F₁- progeny and increasing mortality percentage.

Keywords: Magnetic Field – *Sitophilus oryzae*- *Rizopertha dominica*- Germination

INTRODUCTION

According to a 1990 survey of extension specialists through-out the United States, stored grain losses exceeded \$500 million for the year. Most of these losses resulted from infestation by several species of insect pests and damage by numerous molds and mycotoxins. Losses resulting from insect infestations are wide spread and involve more than loss of quality (Phillip and Meronuck, 1995). Damaged kernels are of lighter weight and result in discounts when marketed. Insect infestation also causes a reduction in nutrients in the grain. Controlling insects with insecticides, including fumigants, rather than using preventative methods incurs great cost. In addition, infestation generally results in dissatisfied customers and related marketing problems that develop from a poor reputation in marketing channels. Wheat containing 32 or more IDK(insect damaged kernels)per 100 grams would result in the wheat being designated as sample grade. Restricting the sale of wheat for livestock feed is a significant loss-a loss that some sellers attempted to reduce by claiming the damage occurred in shipment and should be covered by insurance. This claim is not justified since this type of damage (primarily adult insect emergence holes) could not occur in the short shipment period (7 to 14 days). The insects producing IDK damage require 30 to 45 days for development and emergence from the kernels

Investigations on the influence of magnetic fields on seeds and plants over many years suggest that they lead to better plant growth and yield than chemical fertilizers and contributed to the improvement of the crop productivity and protection. In addition, there have been developed magnetic technologies in several

countries that are ecologically friendly and non-polluting to the soil and are potentially attractive as being affordable to farmers (Liboff *et al.*, 1992; Katsen *et al.*, 2003).

Magnetic Fields (MFs) is not expensive, and at the same time not dangerous to the environment, Other than many radiation sources. (MFs) has attracted the attention of researchers due to their biological effects. Most of the studies about MFs' effects have focused on vertebrates and relatively fewer studies have been done on insects and their stored-product environment (Starick *et al.*, 2005). MFs have been shown to affect the orientation (Jones and Macfadden, 1982), oviposition and development (Ramirez *et al.*, 1983), fecundity and behavior (Starick *et al.*, 2005) of a wide variety of insects.

MATERIALS AND METHODS

The present research was conducted at Seed Technology Research Unit, Field Crops Research Institute and Plant Protection Research Institute - Agriculture Research Centre, Mansoura, Egypt to evaluate the effect of magnetic field on seed germination characters of three wheat varieties (Masr-1, Sids-12 and Sakha-93) infected with two primary stored grain insects rice weevil, *Sitophilus oryzae* (L.) and lesser grain borer, *Rizopertha dominica* F. Seeds were obtained from Wheat Research Department, Field Crops Research Institute, Agriculture Research Centre, Giza, Egypt and were infected with the two insect pests at Plant Protection Research Institute.

Insect rearing:

Adults of *S.oryzae* and *R.dominica* were reared in glass jars (each of approximately 500 ml) containing about 250 gm of wheat seeds. Each jar was covered

with muslin cloths and fixed with rubber bands for egg laying to obtain large numbers of adults needed for the tests and incubated at 30±2°C and 65±5 % R.H.

Bioassay tests:

Twenty adults of *S. oryzae* and *R. dominica* (1-2 week old) infesting 10 gm of three wheat seeds varieties ((Masr-1, Sids-12 and Sakha-93) were exposed to round permanent magnets of about 30 and 60 mT through different time (1 min., 6 and 12 hrs). Each treatment replicated 3 times. After exposure period, the mortality percentage was recorded after 2,4,6,8 and 10 days then the live insects were removed and the replicates were incubated at 30±2°C and 65±5 % R.H. to investigate the reduction in F1-progeny after 60 days.

Germination tests:

Seeds immersed in 5% NaOCl (Sodium hypochloride solution) for 5 min to avoid fungal invasion. Germination tests were performed according to ISTA, (1999), 8 replicates of 50 seeds from each treatment (400 seeds) were placed in Petri dishes (12cm) containing 3 layers of moistened blotters and incubated in the growth chamber at 20± 2°C to study the following parameters:

- Germination percentage defined as the total number of normal seedlings at the end of the test after seven days.

- **Germination rate (GR):** It was calculated according to Bartlett, (1937):

$$GR = \frac{a + (a + b) + (a + b + c) + \dots + (a + b + c + m)}{n(a + b + c + m)}$$

In which a, b, c are No. of seedlings in the first, second and third count, m is No. of seedlings in final count, n is the number of counts.

- **Mean Germination Time (MGT):** calculated based on the equation of Ellis and Roberts (1981). $MGT = \frac{\sum Dn}{\sum n}$

Where (n) is the number of seeds, which were germinated on day, D is number of days counted from the beginning of germination.

- **Speed Germination Index (SGI):** calculated according to the Association of Official Seed Analysis (AOSA., 1983) equation:

$$SGI = \frac{\frac{\text{No. of germinated seed}}{\text{Days of first count}} + \frac{\text{No. of sgerminated seed}}{\text{Days of final count}}}{2}$$

The seeds were considered germinated when the radical was 2 mm long at least

- **Co-efficient of germination (CG):** It was calculated using the following formula (Copeland 1976).

$$\text{Co-efficient of germination} = \frac{100(A1 + A2 + \dots + An)}{A1T1 + A2T2 + \dots + AnTn}$$

Where,

A = Number of seed germinated.

T = Time (days) corresponding to A.

n = No. of days to final count.

- **Seedlings length (cm):** It was measured of ten normal seedlings at 7 days after planting.

- **Seedlings dry weight (gm):** Ten normal seedlings at 7 days after planting, the seedlings were dried in hot-air oven at 85° C for 12 hours to obtain the seedlings dry weight (g).

- **Seedling vigor:** It was calculated according to Abdul Baki and Anderson (1973) as;

$$\text{Vigor index I} = \text{Germination (\%)} \times \text{Seedling length (Root + Shoot)}$$

$$\text{Vigor index II} = \text{Germination (\%)} \times \text{Seedling dry weight (Root + Shoot)}$$

- **Electrical conductivity** of seed leakages was determined for 50 individually weighed seeds per sample placed into 250 mL distilled water at 20°C. After 24 h the leakage electrical conductivity was measured using the CMD 830 WPA (Matthews and Powell, 1981). The conductivity was expressed as µmhos/g seed

Seed Components Analysis:

Moisture %, crude fats%, crude protein%, ash % and carbohydrates % were determined in grains after complete application of magnetic treatments.

Total ash content: 2 gram of grains sample were added into previously weighed porcelain crucible, place in muffle furnace for 2 hours at 600°C then placed in desiccators and weigh. The weight of the residue was calculated and expressed as percent ash (AOAC, 2000).

Crude Fat (Ether Extract): 10 gram of each powdered grains sample were extracted using (Soxhlet) with a solvent of petroleum ether (b.p.60-80°C) for 16 hours. Each extract was dried over anhydrous Na₂SO₄ and evaporated to dryness. The residue was dried at 80°C for 10 minutes, cooled, weighed and expressed as percent lipid (AOAC, 2000).

Crude Fiber Contents: 2 gram of each sample defatted powder were boiled with 1.25% sulphuric acid for 30 minutes and filtered. The residue was dried at 100°C to constant weight the difference between the weight of residue after drying at 110°C and the of powder represents the weight of crude fiber (AOAC, 2000)

Crude Protein: calculated by multiplying the total nitrogen by the factor 6.25. According to (AOAC, 2000),

Moisture Contents: Five g air-dried powder sample were accurately weighed and dried in an oven until constant weight was obtained. The loss in weight was calculated according to (AOAC, 2000)

Determination of total carbohydrate: Carbohydrate percentage was given by: 100 – (percentage of ash + percentage of moisture + percentage of fat + percentage of protein. (Shumaila and Mahpara, 2009).

Statistical analysis

Data were statically analyzed using an analysis of variance (ANOVA) of completely randomized design (MSTAT-C v. 3.1., 1988). Least Significant difference (LSD) was applied to compare mean values.

RESULTS

The adult mortality percentage of *Sitophilus oryzae* and *Rizopertha dominica* after using two MFs levels (30 & 60mT) through different time (1 min., 6 and 12 hrs) are shown in tables (1&2). The mortality percentage of both insects increased with increasing of MFs levels and time. MFs level at 60 mT for 12 hrs was the most effective against both tested insects. *R.dominica* was more resistance than *S.oryzae* in which

56.6, 53.3 and 50.0 % of *S. oryzae* died with Masr-1, Sids-12 and Sakha-93 respectively after 10 days from exposure period while at the same level and time, mortality percentage of *R. dominica* was 45.0, 36.6 and 33.3% with Sids-12, sakha-93 and Masr-1, respectively.

Although, both MFs levels didn't cause high mortality percentage with both insects but they affected on the insects feeding, mating and female fertility leading to a good reduction percentage in F₁-progeny.

Reduction percentage in F₁-progeny of *S. oryzae* ranged from (36.7 – 62.0 %), (30.4 - 41.0%) and (20.5 -36.0 %) at 30 mT level with Masr-1,Sakha-93 and Sids-12 varieties, respectively and at 60 mT level reduction was (36.0 –78.0 %), (37.0 – 74.0 %) and (24.6 -59.0 %) while reduction in F₁-progeny of *R. dominica* ranged from (4.0 – 38.0 %), (29.0 – 59.0 %) and (17.0 – 52.0 %) at 30 mT and at 60 mT reduction (%) was (24.0 - 52.0 %), (37.0 – 63.0 %) and (30.0 – 61.0 %).

Table 1. Effect of magnetic field treatments on *S. oryzae* infesting wheat cultivars and reduction percentage of F₁-progeny at laboratory conditions of 30±2°C and 65±5% R.H.

Treatment Variety	Magnetic	(%) Adult mortality after indicated days					% reduction in F ₁ progeny
		2	4	6	8	10	
Masr1	C ₁	0.00	0.00	0.00	0.00	0.00	
	T ₂	0.0 ± 0.0	1.6 ± 1.6	5.0 ± 2.9	5.0 ± 2.9	5.0 ± 2.9	36.7
	T ₃	6.6 ± 3.3	13.3 ± 3.3	16.6 ± 4.4	21.6 ± 3.3	28.3 ± 4.4	41.0
	T ₄	11.6 ± 4.4	20.0 ± 2.9	28.3 ± 6.0	30.0 ± 5.7	36.6 ± 5.8	62.0
	T ₅	0.0 ± 0.0	3.3 ± 3.3	8.3 ± 4.4	10.0 ± 2.9	10.0 ± 2.9	36.0
	T ₆	8.3 ± 3.3	13.3 ± 1.7	21.6 ± 1.7	33.3 ± 5.8	35.0 ± 4.9	75.0
	T ₇	15.0 ± 5.7	25.0 ± 5.7	36.6 ± 6.0	43.3 ± 4.4	56.6 ± 8.6	78.0
Sids12	C ₁	0.00	0.00	0.00	0.00	0.00	
	T ₂	1.6 ± 1.6	3.3 ± 1.6	5.0 ± 2.8	5.0 ± 2.8	5.0 ± 2.8	30.4
	T ₃	1.6 ± 1.6	5.0 ± 2.7	11.6 ± 1.6	15.0 ± 2.8	16.6 ± 3.3	33.0
	T ₄	3.3 ± 1.6	10.0 ± 2.8	18.3 ± 1.6	21.6 ± 4.4	23.3 ± 3.3	41.0
	T ₅	1.6 ± 1.6	6.6 ± 3.3	8.3 ± 1.6	10 ± 2.8	11.6 ± 1.6	37.0
	T ₆	3.3 ± 1.6	10 ± 5	15.0 ± 2.8	23.3 ± 3.3	30.0 ± 2.8	61.0
	T ₇	16.6 ± 1.6	28.3 ± 1.6	38.3 ± 6.0	46.6 ± 10.0	50.0 ± 8.6	74.0
Sakha93	C ₁	0.00	0.00	0.00	0.00	0.00	
	T ₂	0.0 ± 0.0	0.0 ± 0.0	1.6 ± 1.6	5.0 ± 2.8	5.0 ± 2.8	20.5
	T ₃	1.6 ± 1.6	1.6 ± 1.6	5.0 ± 4.9	8.3 ± 3.3	10.0 ± 2.8	34.0
	T ₄	23.3 ± 7.2	33.3 ± 4.4	35 ± 4.9	43.3 ± 9.2	46.6 ± 8.7	36.0
	T ₅	0.0 ± 0.0	0.0 ± 0.0	3.3 ± 3.3	5.0 ± 2.8	6.6 ± 1.6	24.6
	T ₆	3.3 ± 1.6	6.6 ± 1.6	10.0 ± 4.9	13.3 ± 3.3	20.0 ± 2.8	46.5
	T ₇	31.6 ± 4.4	35 ± 10.4	40.0 ± 8.6	48.3 ± 8.6	53.3 ± 10.9	59.0
F.sig.		**	**	ns	ns	ns	
LSD at 5%		1.61	2.00	ns	ns	ns	
Variety	Misar-1	5.9 ± 1.6	10.9 ± 2.2	16.6 ± 3.3	20.5 ± 3.5	24.5 ± 4.5	
	Sides-12	4 ± 1.3	9 ± 2.1	13.8 ± 2.7	17.4 ± 3.5	19.5 ± 3.7	
	Sakha-93	8.6 ± 2.9	10.9 ± 3.6	13.6 ± 3.8	17.6 ± 4.4	20.2 ± 4.5	
F.sig.		*	ns	ns	Ns	ns	
LSD at 5%		0.61	ns	ns	Ns	ns	
Magnetic treatment	C ₁	0.00	0.00	0.00	0.00	0.00	
	T ₂	0.6 ± 0.6	1.6 ± 0.8	3.9 ± 1.4	5 ± 1.4	5 ± 1.4	
	T ₃	3.3 ± 1.4	6.7 ± 2.2	11.1 ± 2.6	15 ± 2.5	18.3 ± 3.2	
	T ₄	12.7 ± 3.8	21.1 ± 3.8	27.2 ± 3.3	31.7 ± 4.6	35.6 ± 4.7	
	T ₅	0.6 ± 0.6	3.3 ± 1.6	6.7 ± 1.9	8.3 ± 1.7	9.4 ± 1.3	
	T ₆	5 ± 1.4	10 ± 1.8	15.6 ± 2.4	23.3 ± 3.6	28.3 ± 2.9	
	T ₇	21.1 ± 3.4	29.4 ± 3.8	38.3 ± 3.5	46.1 ± 4.1	53.3 ± 4.8	
F.sig.		***	***	***	***	***	
LSD at 5%		00.93	1.16	1.33	1.57	1.63	
CV (%)		79.48	58.20	47.97	44.68	40.09	

C₁ : Control without infection without magnetic T₂ : Expose infected seeds with 30 mT for 1 minuet.
 T₃ : Expose infected seeds with 30 mT for 6 hours. T₄ : Expose infected seeds with 30 mT for 12 hours.
 T₅ : Expose infected seeds with 60 mT for 1 minuet. T₆ : Expose infected seeds with 60 mT for 6 hours.
 T₇ : Expose infected seeds with 60 mT for 12 hours. CV : Coefficient of variation

Table 2. Effect of magnetic field treatments on *R. dominica* infesting wheat cultivars and reduction percentage of F₁-progeny at laboratory conditions 30±2°C and 65±5% R.H.

Treatment	(%) Adult mortality after indicated days						% reduction in
Variety	Magnetic	2	4	6	8	10	F ₁ progeny
Masr-1	C ₁	0.00	0.00	0.00	0.00	0.00	
	T ₂	0 ± 0	0 ± 0	0 ± 0	3.3 ± 1.6	3.3 ± 1.6	17.0
	T ₃	0 ± 0	0 ± 0	3.3 ± 1.6	6.6 ± 4.4	6.6 ± 4.4	48.0
	T ₄	3.3 ± 3.3	6.6 ± 3.3	8.3 ± 1.6	10 ± 0	13.3 ± 1.6	52.0
	T ₅	5 ± 2.8	6.6 ± 1.6	11.6 ± 1.6	11.6 ± 1.6	11.6 ± 1.6	30.0
	T ₆	6.6 ± 4.4	13.3 ± 4.4	15 ± 2.8	16.6 ± 3.3	18.3 ± 1.6	43.0
	T ₇	6.6 ± 3.3	15 ± 2.8	21.6 ± 3.3	26.6 ± 4.4	33.3 ± 4.4	61.0
Sids-12	C ₁	0.00	0.00	0.00	0.00	0.00	
	T ₂	0 ± 0	3.3 ± 1.6	5 ± 0	8.3 ± 1.6	8.3 ± 1.6	29.0
	T ₃	0 ± 0	5 ± 2.8	8.3 ± 1.6	11.6 ± 4.4	11.6 ± 4.4	44.0
	T ₄	5 ± 2.8	8.3 ± 1.6	15 ± 2.8	20 ± 2.8	20 ± 2.8	59.0
	T ₅	1.6 ± 1.6	11.6 ± 4.4	11.6 ± 4.4	11.6 ± 4.4	11.6 ± 4.4	37.0
	T ₆	15 ± 2.8	23.3 ± 4.4	23.3 ± 4.4	26.6 ± 7.2	26.6 ± 7.2	45.0
	T ₇	21.6 ± 1.6	31.6 ± 6	36.6 ± 3.3	41.6 ± 1.6	45 ± 2.8	63.0
Sakha-93	C ₁	0.00	0.00	0.00	0.00	0.00	
	T ₂	0 ± 0	0 ± 0	3.3 ± 1.6	3.3 ± 1.6	3.3 ± 1.6	4.0
	T ₃	0 ± 0	0 ± 0	5 ± 2.8	10 ± 4.4	10 ± 2.8	19.0
	T ₄	1.6 ± 1.6	8.3 ± 1.6	11.6 ± 1.6	15 ± 4.9	18.3 ± 7.2	38.0
	T ₅	1.6 ± 1.6	6.6 ± 3.3	10 ± 5.7	13.3 ± 4.4	13.3 ± 4.4	24.0
	T ₆	5 ± 2.8	13.3 ± 3.3	16.6 ± 3.3	16.6 ± 3.3	16.6 ± 3.3	43.0
	T ₇	11.6 ± 4.4	20 ± 5.7	26.6 ± 3.3	36.6 ± 3.3	36.6 ± 3.3	52.0
F.sig.		*	ns	ns	ns	ns	
LSD at 5%		1.21	ns	ns	ns	ns	
Variety	Misar-1	3.1 ± 1.1	6 ± 1.5	8.6 ± 1.8	10.7 ± 2	12.4 ± 2.5	
	Sides-12	6.2 ± 1.9	12 ± 2.6	14.3 ± 2.7	17.1 ± 3.1	17.6 ± 3.3	
	Sakha-93	2.8 ± 1.1	7 ± 1.8	10.5 ± 2.1	13.6 ± 2.7	14 ± 2.7	
F.sig.		*	**	**	**	*	
LSD at 5%		0.46	0.64	0.59	0.72	0.77	
Magnetic treatment	C ₁	0.00	0.00	0.00	0.00	0.00	
	T ₂	0.0 ± 0	1.1 ± 0.7	2.7 ± 0.9	5 ± 1.2	5 ± 1.2	
	T ₃	0.0 ± 0	1.7 ± 1.2	5.6 ± 1.3	9.4 ± 2.1	9.4 ± 2.1	
	T ₄	3.3 ± 1.4	7.8 ± 1.2	11.7 ± 1.4	15 ± 2.2	15 ± 2.2	
	T ₅	2.8 ± 1.2	8.3 ± 1.8	11.1 ± 2.2	12.2 ± 1.9	12.2 ± 1.9	
	T ₆	8.9 ± 2.3	16.6 ± 2.6	18.3 ± 2.2	20 ± 3	20.6 ± 2.8	
	T ₇	13.3 ± 2.8	22.2 ± 3.5	28.3 ± 2.8	35 ± 2.8	38.3 ± 2.5	
F.sig.		***	***	***	***	***	
LSD at 5%		0.70	0.98	0.90	1.11	1.17	
CV (%)		95.9	62.93	42.8	42.30	42.04	

*Explanations are as in Table 1

Seed vigor traits i.e. germination rate (GR), mean germination time (MGT), speed germination index (SGI) and co-efficient of germination (CG) as affected by varieties and magnetic treatments are shown in table 3. The tested three wheat varieties (Masr-1, Sids-12 and Sakha-93) showed equal response in all seed vigor traits under infection with *R. dominica*. While slightly differences were recorded in seed vigor traits except SGI under infection with *S. oryzae*.

Regarding magnetic treatments, data in the same table showed that exposure infected seeds with (*S. oryzae* or *R. dominica*) to magnetic field treatments positively affected on all seed vigor tested parameters compared control treatment (infected seeds without magnetic treatment; C₂). Moreover, almost equaled with uninfected seeds (C₁). Expose infected seeds with both

insects at 60 mT increased gradually with increasing exposes time from 1 min to 6 and 12 hrs in all tested parameters except MGT where reverse trend was recorded. Generally, T₇ recorded the highest value without significant with T₆ in the most tested parameters under both insects. The percentage of improvements regarding application of different magnetic treatments compared to control (C₂) ranged between 10.68-17.95%, 15.02-24.82%, 16.22-20.89% and 17.57 - 33.08% in GR, MGT, SGI and CG, respectively under seeds infected with *R. dominica*. Similar trends with higher value were recorded under *S. Oryzae* where the increasing percent ranged between 14.68-20.63%, 19.94-27.68%, 28.59-33.18% and 22.73-35.18% in the above mentioned parameters, respectively.

Regarding the interactions between varieties and magnetic treatments, results in Table 4, showed

insignificant effects on all tested parameters except GR, MGT and CG under infected seeds with *S. oryzae*. Generally, application of different magnetic treatments

on the tested three varieties improved all tested parameters compared with infected seed without magnetic treatment under both insects.

Table 3. Effect of magnetic field treatments on germination rate (GR), mean germination time (MGT), speed germination index (SGI) and co-efficient of germination (CG) of three wheat varieties infected with *R. dominica* or *S. oryzae*.

Character		<i>R. dominica</i>				<i>S. oryzae</i>			
Treatment		GR	MGT	SGI	CG	GR	MGT	SGI	CG
Variety	Misar-1	0.88	1.37	20.88	74.35	0.90	1.29	20.98	78.07
	Sides-12	0.88	1.37	20.74	74.25	0.86	1.42	20.30	72.61
	Sakha-93	0.87	1.39	20.64	73.46	0.89	1.34	20.90	76.80
F test		ns	Ns	Ns	Ns	**	**	ns	**
LSD at 5%		ns	Ns	Ns	Ns	0.02	0.06	ns	3.50
Magnetic seeds	#C ₁	0.94	1.19	22.33	84.17	0.94	1.19	22.11	84.00
	C ₂	0.77	1.68	17.81	59.66	0.77	1.70	16.52	60.43
	T ₅	0.86	1.43	20.70	70.15	0.88	1.36	21.24	74.16
	T ₆	0.90	1.31	21.54	76.73	0.91	1.28	21.76	78.85
	T ₇	0.91	1.26	21.39	79.40	0.92	1.23	22.00	81.69
F test		**	**	**	**	**	**	**	**
LSD at 5%		0.03	0.07	0.75	3.43	0.03	0.08	1.19	4.52
CV (%)		2.56	4.89	3.77	4.82	3.25	6.41	5.96	6.20

C₁ : Control without infection without magnetic C₂: Control with infection without magnetic
 T₅ : Expose infected seeds with 60 mT for 1 minute. T₆ : Expose infected seeds with 60 mT for 6 hours.
 T₇ : Expose infected seeds with 60 mT for 12 hours. CV: Coefficient of variation

Table 4. Effect of magnetic field treatments on germination rate (GR), mean germination time (MGT), speed germination index (SGI) and co-efficient of germination (CG) of three wheat varieties infected with *R. dominica* or *S. oryzae*.

Character		<i>R. dominica</i>				<i>S. oryzae</i>			
Treatment	Magnetic	GR	MGT	SGI	CG	GR	MGT	SGI	CG
Variety	#C ₁	0.93	1.20	22.50	83.33	0.94	1.19	21.67	83.74
		0.78	1.66	18.17	60.31	0.87	1.39	18.17	73.06
		0.85	1.44	20.44	69.70	0.90	1.31	21.61	76.73
		0.90	1.31	21.61	76.39	0.89	1.32	21.44	76.11
		0.93	1.22	21.67	82.03	0.92	1.24	22.00	80.70
Misar-1	C ₂	0.94	1.17	22.00	85.84	0.94	1.18	22.17	84.92
		0.78	1.66	17.78	60.37	0.71	1.86	15.56	53.64
		0.86	1.43	21.33	70.11	0.84	1.47	20.61	67.94
		0.91	1.27	21.56	78.98	0.90	1.30	21.33	77.11
		0.89	1.32	21.06	75.95	0.91	1.26	21.83	79.44
Sides-12	T ₅	0.93	1.20	22.50	83.33	0.93	1.20	22.50	83.33
		0.76	1.72	17.50	58.31	0.72	1.83	15.83	54.59
		0.86	1.42	20.33	70.64	0.90	1.29	21.50	77.83
		0.88	1.35	21.44	74.81	0.93	1.20	22.50	83.33
		0.92	1.25	21.44	80.22	0.94	1.18	22.17	84.92
Sakha-93	T ₆	ns	Ns	ns	Ns	**	**	ns	**
		ns	Ns	ns	Ns	0.05	0.14	ns	7.83

#Explanations are as in Table 3

Response of three wheat varieties infected with *S. oryzae* or *R. dominica* to exposing of different magnetic treatments on changes in germination%, seedling length (cm), seedling dry weight (g), seedling vigor index (SVI) and electrical conductivity (EC) of seeds are shown in Table 5 and 6.

The tested three wheat varieties differed slightly for mentioned characters under infection with both tested insects where Masr-1 gave the highest value of SL, SDW and SVI, followed by Sides-12 and Sakha-93,

respectively. While under infection with *S. oryzae*, the order was Sakha-93, Sides-12 and Maser-1, respectively (Table 5).

Regarding magnetic treatments, results in the same Table showed that expose the infected seeds to magnetic field 60 mT for 1 min, 6 and 12 hrs caused positive significant effects on all tested characters compared to infected seeds without magnetic treatment; (C₂). The percent of improvement ranged between 2.08 – 3.47%, 14.19 – 21.34%, 22.43 – 44.45% and 25.08 –

49.75% in G %, SL, SDW, and SVI, respectively under infected with *R. dominica*. Whereas, magnetic treatments after infection with *S. oryzae*, the increasing percent ranged between 2.82 – 4.23%, 10.79 – 22.35%, 55.19 – 93.34%, 58.93 – 100.70% in the above mentioned parameters, respectively.

Results in Table (6) show significant effects regarding the interactions between wheat varieties and magnetic field treatments on seedling dry weight and SVI under infected seed with *R. dominica* and *S. oryzae*.

While no significant effects were reported in germination % and seedling length under both insects. Generally, Sakha-93 came in the first order followed by Masar-1 and sides-12 for producing the heaviest seedling dry weight and highest value of SVI under different treatments.

Electrical conductivity result showed that high significant effect by using mf, the lowest values resulted from exposing wheat seed to 60 mT for 12 h in all infected varieties; these results refer to high seed vigor

Table 5.Effect of magnetic field treatments on germination %, seedling length (SL), seedling dry weight (SDW), seedling vigor index (SVI) and electrical conductivity (EC) of three wheat varieties infected with *R. dominica* or *S. oryzae*.

Character	Treatment	<i>R. dominica</i>					<i>S. oryzae</i>				
		Germ. (%)	Seedling length (cm)	Seedling dry wt. (g)	SVI	EC	Germ. (%)	Seedling length (cm)	Seedling dry wt. (g)	SVI	EC
Variety	Misar-1	99.60	22.70	0.132	13.20	0.015	97.20	21.179	0.370	35.85	0.011
	Sides-12	97.20	22.53	0.117	11.40	0.012	98.00	22.756	0.390	38.37	0.011
	Sakha-93	98.80	20.86	0.145	14.38	0.010	98.40	20.766	0.480	47.81	0.013
F test		**	**	**	**	**	ns	**	**	**	**
LSD at 5%		1.39	0.868	0.01	0.86	0.003	ns	1.011	0.020	2.97	0.001
Magnetic seeds	#C ₁	100.00	24.18	0.156	15.63	0.009	100.00	23.95	0.547	54.72	0.009
	C ₂	96.00	18.99	0.101	9.65	0.019	94.67	18.56	0.248	23.56	0.017
	T ₅	98.00	21.68	0.123	12.07	0.013	97.33	20.56	0.385	37.45	0.011
	T ₆	99.33	22.25	0.133	13.17	0.011	98.67	22.07	0.410	40.37	0.010
	T ₇	99.33	23.04	0.145	14.45	0.010	98.67	22.70	0.479	47.29	0.010
F test		**	**	**	**	**	**	**	**	**	**
LSD at 5%		1.79	1.121	0.01	1.11	0.004	2.98	1.306	0.03	3.83	0.002
CV (%)		1.89	5.25	9.04	8.84	23.65	3.17	6.290	9.24	9.78	15.44

#Explanations are as in Table 3

Table 6.Effect of magnetic field treatments on germination %, seedling length (SL), seedling dry weight (SDW), seedling vigor index and electrical conductivity (EC) of three wheat varieties infected with *R. dominica* or *S. oryzae*.

Character	Treatment	<i>R. dominica</i>					<i>S. oryzae</i>					
		Magnetic	Germ. (%)	Seedling length (cm)	Seedling dry wt. (g)	SVI	EC	Germ. (%)	Seedling length (cm)	Seedling dry wt. (g)	SVI	EC
Misar-1	#C ₁		100.00	25.38	0.154	15.43	0.009	100.00	24.50	0.447	44.68	0.009
	C ₂		98.00	19.45	0.093	9.11	0.024	90.00	17.11	0.232	20.90	0.020
	T ₅		100.00	21.99	0.132	13.23	0.014	96.00	19.14	0.342	32.83	0.009
	T ₆		100.00	22.79	0.133	13.30	0.014	100.00	22.43	0.375	37.53	0.009
	T ₇		100.00	23.88	0.149	14.93	0.013	100.00	22.72	0.433	43.32	0.009
Sides-12	#C ₁		100.00	25.11	0.136	13.57	0.009	100.00	25.75	0.518	51.77	0.009
	C ₂		94.00	19.49	0.104	9.81	0.018	96.00	20.01	0.186	17.84	0.014
	T ₅		96.00	22.24	0.107	10.21	0.014	98.00	22.38	0.383	37.47	0.011
	T ₆		98.00	22.38	0.119	11.62	0.009	98.00	22.41	0.401	39.33	0.009
Sakha-93	T ₇		98.00	23.42	0.120	11.77	0.009	98.00	23.23	0.463	45.42	0.009
	#C ₁		100.00	22.06	0.179	17.88	0.009	100.00	21.59	0.677	67.71	0.010
	C ₂		96.00	18.03	0.105	10.02	0.014	98.00	18.55	0.325	31.94	0.018
	T ₅		98.00	20.81	0.131	12.77	0.010	98.00	20.16	0.429	42.03	0.013
	T ₆		100.00	21.59	0.146	14.59	0.009	98.00	21.36	0.452	44.24	0.011
T ₇		100.00	21.82	0.167	16.65	0.009	98.00	22.16	0.541	53.12	0.011	
F test		Ns	ns	*	*	ns	ns	ns	**	*	*	
LSD at 5%		Ns	ns	0.02	1.91	ns	ns	ns	0.05	6.63	0.003	

#Explanations are as in Table 3

Seed Components Analysis:

Means of three wheat varieties seeds components including moisture content %, crude fats%, crude protein%, ash % and carbohydrates % after using MF level 60mT for 12 hrs are shown in tables 7. Control seeds were used for comparison.

Results showed that, MF level slightly increase moisture content (%) and total carbohydrate (%) while slightly decrease crude protein (%), total fats (%) and crude fiber (%) after 12 hrs exposure. Although, increasing of seed moisture content, still under limited for safe storage. Prior of storage, the moisture content of

cultivars seed as good quality based on certification standard cited by (Copeland and McDonald 2004) have recommendation which moisture content for long duration storage seed does not above 14 or below 5 %. Seeds store at moisture content above 14% begin to exhibit increased respiration, heating and fungal invasion that destroy seed viability more rapidly, while below 5 % cause seed membrane structure , seed deterioration, thus improve that it had no significant effect on seeds viability

Table 7. Effect of static magnetic field at 60 mT for 12hrs on grains chemical constituents (%) of three wheat varieties infested with *S.oryzae* and *R.dominica*.

Character	Treatment	Grains chemical constituents (%)					
		Moisture	Crude protein	Total fat	Ash	Total Carbohydrates	Crude fiber
Masr-1	Control	10.30	11.78	2.40	2.27	73.52	2.58
	<i>S.oryzae</i>	11.63	8.84	1.74	1.66	74.55	1.85
	<i>R.dominica</i>	12.95	9.63	0.98	1.05	75.39	1.14
Sids-12	Control	9.16	12.47	2.87	2.66	72.84	3.05
	<i>S.oryzae</i>	10.49	11.41	2.16	2.06	73.88	2.30
	<i>R.dominica</i>	11.84	10.33	1.49	1.45	74.89	1.61
Sakha-93	Control	9.61	12.1	2.62	2.48	73.19	2.81
	<i>S.oryzae</i>	10.92	11.04	1.98	1.84	74.22	2.07
	<i>R.dominica</i>	12.59	9.97	1.25	1.27	74.92	1.36

DISCUSSION

The development of alternative treatments for pest control is an increasing demand from the food industry. Alternatives should meet consumer demands for the reduced use or elimination of pesticides, while at the same time maintaining a high degree of control efficacy (Riudavetset *al.*, 2010). The present study established the efficacy of using magnetic fields (MFs) to control stored product pests. The advantages of magnetization as a pest control methods that it does not leave undesirable residues. Other few researchers studied the biological effect of MFs on stored grain insects like (Pandiret *al.*, 2013a) who investigate the effect of MFs from a DC power supply on the longevity of adults, fecundity and daily egg laying pattern of female *Ephestia kuehniella* (Zeller) and found that mortality increased with increasing MFs level and complete mortality was achieved at the level of 10 mT. Pest longevity was significantly reduced with increasing level of MFs. There was no significant difference in the longevity of males and females. Exposing adults to increasing level of MFs significantly influenced daily egg laying patterns and fecundity of magnetized females. Larval emergence from these eggs was completely prevented at highest level of MFs. (Pandir and Sahingoz 2014) investigation showed that MFs caused oxidative stress and proved to be DNA damage as revealed by the comet assay. MFs may be used to determine potential toxic effects as a control agent against *E. kuehniella* larvae. (Pandiret *al.*, 2013b). Investigated the effect of strong magnetic fields

as insecticidal activity on *E.kuehniella* (Lepidoptera: Pyralidae) larvae and eggs at different stages of development and their preference by the egg parasitoid, *Trichogramma embryophagum*. Eggs ranging in age from 24-h to 48-h and 72-h-old and larvae (1 to 2 days) were exposed to 1.4 Tesla (T) magnetic fields from a DC power supply at 50 Hz for different time periods (3, 6, 12, 24, 48 and 72 h). Twelve hours of exposure at 1.4 T was toxic to 24-h-old eggs of *E. kuehniella*. The 72-h-old host eggs treated with 1.4 T for 6-72 h were not significantly preferred by *T. embryophagum*. The magnetic field was toxic to 24-h-old eggs of *E. kuehniella* exposed to 1.4 T for 12. The treatment of magnetic fields on the 72-h-old host egg with 1.4 T at 6-72 h was not significantly preferred by *T. embryophagum*. Magnetization of 24-h-old eggs of *E. kuehniella* for 3 h could be effectively used with *T. embryophagum* as sterilised host eggs. These eggs were markedly preferred by *T.embryophagum*. The LT_{50} and LT_{99} values of magnetic fields at different egg stages of *E. kuehniella*, and larvae were measured. A level of 1.4 T at 72 h completely prevented the development of the larvae. There was no significant effect on larval survival at 1.4 T at 48 and 72 h. Increasing magnetic fields exposure times for eggs that were 24-h, 48-h and 72-h-old prevented larval emergence and increased their mortality rate. Consequently, magnetic fields could be used in controlling stored-product pest eggs and larvae of *E. kuehniella*.

Results showed that exposure infected seeds with (*Sitophilus oryzae* and *Rizopertha dominica*) to

magnetic field treatments positively affected on all seed and seedling vigor parameters compared control treatment (infected seed without magnetic treatment; C₂). Moreover almost equaled with uninfected seeds but it showed significant effect on all parameters comparing with untreated seeds (C₁). MF affected the various characteristics of the plants like germination of seeds, root growth, rate seedlings growth, reproduction and growth of the meristem cells and chlorophyll quantities (Namba *et al.*, 1995; Atak *et al.*, 1997; Reina *et al.*, 2001; Atak *et al.*, 2003 and Aycih and Alikamanoglu 2005) concluded that magnetic field increased the shoot and root regeneration rate and their fresh weight in soybean and paulownia organ cultures. (Pouya and Nasrin 2015) found that, an application of alternating magnetic field of (B=0.5mT and f=50Hz) on seed affected seedling growth. Longer exposure time (40 s) has generally induced slightly growth. The resulting differences in fresh weight of seedling were not correlated with the differences in shoot heights, In 20 seconds period, the specific electromagnetic field intensity, increased the stimulation of wheat buds and gained germination percentage rather to the control group, but shifting the effective time to 40 seconds, the growth decreased to some extent. (Cakmak *et al.*, 2010) observed that the application of magnetic field doses of 4 mT and 7 mT promoted germination ratios of bean and wheat seeds. (De Souza *et al.*, 2010) concluded that pre-sowing magnetic treatments have the potential to enhance tomato seed germination and early seedling growth. Alvarez *et al.*, (2012) concluded that magnetic treatment improves germination rate of triticale seeds. In general, most of the parameters recorded for all the doses applied to triticale seeds were better than control values. Furthermore, seedlings from magnetically treated seeds grew taller than control.

REFERENCES

- Abdul-Baki, A.A. and J.D. Anderson (1973) Vigour determination in soybean by multiple criteria. *Crop Sci.* 10:31-34.
- Alvarez, J.; M.V. Carbonell ; M. Flórez; E. Martínez and A. Campos (2012) Study of 125 mT magnetic treatment on the germination and initial growth of triticale seeds. *Научни Трудове На Русенския Университет - том 51, серия 3.1*
- AOAC, (2000) Association of Official Analytical Chemists. (17th ED. Of A.O.A.C. international published by A.O.A.C. international Maryland, U.S.A., 1250pp).
- AOSA, (1983) Seed Vigor Testing Handbook: Contribution No. 32 to the Handbook on Seed Testing. Association of Official Seed Analysis (AOSA), Springfield, IL., USA., Pages: 93.
- Atak, C.; O. Emiroglu; S. Aklimanoglu and A. Rzakoulieva (2003). Stimulation of regeneration by magnetic field in soybean (*Glycine max* L. Merrill) tissue cultures. *J Cell Mol. Biol.*, 2:113-119.
- Atak, C.; V. Danilov; B. Yurttas; S. Yalçın; D. Mutlu and A. Rzakoulieva (1997) Effects of magnetic field on soybean (*Glycine max* L. Merrill) seeds. *Com JINR. Dubna*, 1-13.
- Bartlett, M.S. (1937) Some samples of statistical method of research in agriculture and applied biology. *J. Roy. Soc.* 4:2.
- Cakmak, T.; R. Dumlupinar and S. Erdal (2010) Acceleration of germination and early growth of wheat and bean seedlings grown under various magnetic field and osmotic conditions. *Bioelectromagnetics*, 31 (2):120-129.
- Copeland, L.O and M.B. McDonald (2004) Principles of Seed Science and Technology. (4th edition. Kluwer Academic Publishers. Boston, Dordrecht, London).
- Copeland, L.O. (1976). Principles of Seed Science and Technology, Burgess Pub. Com., Minneapolis, Minnesota: 164-165.
- De Souza, A.; L. Sueiro; D. García and E. Porras (2010) Extremely low frequency non-uniform magnetic fields improve tomato seed germination and early seedling growth. *Seed Sci. and Technol.*, 38: 61-72.
- EL-lakwah, F.A.; A.A. Darwish and Z.A. Halawa (1996) Toxic effect of extracts and powders of some plants against the cowpea beetle (*Callosobruchus maculatus*, F.). *Ann. of Agric. Sci.-Moshtohor*, 34 (4): 1849-1859.
- Ellis, H.R.A. and E.H. Roberts (1981) The quantifications of ageing and survival in orthodox seed. *Seed Sci. Technol.* 9:373-407.
- ISTA (1999). International Seed Testing Association. International rules for seed testing. *Seed Sci. Technol. Proc. Int. Seed Test. Ass.* 31(1):1-152.
- Jones, D.S. and B.J. Macfadden (1982) Induced magnetization in the Monarch Butterfly, *Danaus plexippus* (Insecta: Lepidoptera). *J. Exp. Biol.*, 96: 1-9.
- Katsen, A.; T. Dat ; Y. Yogev and A. Prilutsky (2003) Method and Devices for Treatment of a Biological Material with a Magnetic Field. USP atent 6539664 <http://www.freepatentsonline.com/6539664.html>
- Liboff, A.R.; B.R. McLeod and S.D. Smith (1992) Method and Apparatus for Controlling Plant Growth. USP atent 5077934. <http://www.freepatentsonline.com/5077934.html>
- Matthews, S. and A.A. Powell (1981) Electrical Conductivity Test. In: Vigour Test Handbook, Perry, D.A. (Ed.). International Seed Testing Association, Zurich, Switzerland, pp: 37-41.
- MSTAT-C (1988) MSTAT-C, a microcomputer program for the design, arrangement and analysis of agronomic research. Michigan state university, East Lansing.
- Namba, K.; A. Sasao and S. Shibusawa (1995) Effect of magnetic field on germination and plant growth. *Acta Horticulture*. 399: 143-147.

- Pandir, D. and R. Sahingoz (2014) Magnetic field-induced oxidative stress and DNA damage in Mediterranean flour moth *Ephestia kuehniella* Zeller (Lepidoptera: Pyralidae) larvae. J.Pest Sci.; 87(1):79-87. 53 ref.
- Pandir, D.; F. S.Ercan and R. Sahingoz (2013a) Assessment of the influence of magnetic fields on aspects of the biology of the adult Mediterranean flour moth *Ephestiakuehniella* Zeller, 1879 (Lepidoptera: Pyralidae).TurkiyeEntomolojiDergisi;37(4):423-431.
- Pandir, D.; R. Sahingozand F. S. Ercan (2013b) Mediterranean flour moth *Ephestia kuehniella* eggs and larvae exposed to a static magnetic field and preference by *Trichogramma embryophagum*. Biocontrol Science and Technology; 23(12):1402-1411.
- Phillip, H.and R. Meronuck (1995) Stored Grain Losses Due to Insects and Molds and the Importance of Proper Grain Management. J.Sci.Food and Agric.: 29-31
- Pouya, F. and N. Seyedpour (2015).Effects of 50Hz Electromagnetic Fields on Seed Germination and Early Growth in Wheat (*Triticum* spp.). Bull. Env.Pharmacol. Life Sci., Vol 2 (5) April2013:52-54
- Ramirez, E.; J.L. Monteagudo; M. García-Gracia and J.M. Delgado (1983) Oviposition and development of *Drosophila* modified by magnetic fields. Bioelectromagnetics, 4(4): 315-326.
- Renia F.G.;L.APascual.andI.A. Fundora (2001) Influence of a Stationary Magnetic Field on water relations in lettuce Seeds. Part II: Experimental Results. Bioelectromagnetics 22:596-602.
- Riudavets, J.; C. Castañé ; O. Alomar; M.J. Pons and R. Gabarra (2010) The use of carbon dioxide at high pressure to control nine stored-product pests. J. Stored Prod. Res., 46: 228-233.
- Shumaila G. and S. Mahpara (2009) Proximate Composition and Mineral Analysis of Cinnamon. Pakistan Journal of Nutrition, 8(9): 1456-1460.
- Starick, N.T.; B.C. Longstaff and B. Condon (2005) The influence of fluctuating low-level magnetic fields on the fecundity and behavior of *Rizopertha dominica* (F.). J. Stored Prod.t Res., 41: 255-270.

تأثير المجال المغناطيسي على حيوية وجودة البذور والاصابة الحشرية لبعض اصناف القمح محمود حزين^١، دعاء محمد زين العابدين برعي^٢ و أمل على المهدي^٣

^١ قسم بحوث المحاصيل الحقلية- الشعبة الزراعية والبيولوجية - المركز القومي للبحوث
^٢ قسم أفات الحبوب والمواد المخزونة - معهد بحوث وقاية النباتات- مركز البحوث الزراعية - الجيزة
^٣ قسم بحوث تكنولوجيا البذور - معهد محاصيل حقلية - مكن البحوث الزراعية - الجيزة

تم اجراء التجارب المعملية لاختبار فعالية قوتين منخفضتين من المجال المغناطيسي (٣٠ & ٦٠ مللي تسلا) لثلاث فترات مختلفة من التعرض (ادقيقة , ٦ ساعات , ١٢ ساعة) على ثلاث اصناف مختلفة من القمح (مصر ١ , سدس ١٢ , سخا ٩٣) ضد حشرتي سوسة الارز وناقبة الحبوب الصغرى كما تم حساب نسبة الخفض في تعداد الجيل الاول وايضا اختبار تأثير القوتين محل الدراسة على انبات وحيوية البذور وقد اظهرت النتائج:ازداد نسبة الموت بزيادة القوة المستخدمة ومده التعرض . اظهرت حشرة ناقبة الحبوب الصغرى مقاومة اعلى من سوسة الارز مع جميع القوى المستخدمة . سجلت حشرة سوسة الارز نسب موت ٥٦.٦ , ٥٣.٣ و ٥٠ % بعد ١٠ ايام من التعرض مع اصناف مصر ١ , سدس ١٢ وسخا ٩٣ على التوالي بينما سجلت حشرة ناقبة الحبوب الصغرى نسب موت ٤٥ , ٣٦.٦ و ٣٣.٣ مع اصناف سدس ١٢ , سخا ٩٣ و مصر ١ على التوالي مع اعلى قوة مستخدمة ٦٠ مللي تسلا لمدته ١٢ ساعة تعرض . على الرغم من نسب الموت المنخفضة اظهرت النتائج قدرة المجال المغناطيسي (٦٠ مللي تسلا) على خفض تعداد الجيل الاول بنسب عالية وصلت ل ٧٨ % مع سوسة الارز و ٦٣ % مع ناقبة الحبوب الصغرى .واستكمالاً أجريت تجربة انبات بمعامل قسم بحوث تكنولوجيا البذور بالمنصورة لدراسة تأثير المعاملة بالمجال المغناطيسي ٦٠ مللي تسلا لمدة ١ دقيقة , ٦ و ١٢ ساعة على انبات وقوة بادرات اصناف القمح تحت الدراسة (مصر ١ , سدس ١٢ و سخا ٩٣) المصابة بنوعين من حشرات المخازن (سوسة الارز و ناقبة الحبوب الصغرى) . ويمكن تلخيص أهم النتائج فيما يلي اظهرت الاصناف الثلاثة استجابة متساوية في جميع الصفات (معدل الانبات , متوسط زمن الانبات, دليل سرعة الانبات و طاقة الانبات) في حالة الاصابة بحشرة ناقبة الحبوب الصغرى , وباستثناء دليل سرعة الانبات اظهرت اختلافات واضحة لجميع الصفات مع الاصابة بسوسة الارز. تأثرت جميع الصفات بالتعرض للمجال المغناطيسي مقارنة بالكنترول (بذور مصابة بدون معاملة ماجنتيك) وتساوت في معظمها مع الكنترول (الغير معاملة او مصاب) و كانت الزيادة باطالة فترة التعرض للمجال المغناطيسي حيث اعطي التعرض ل ٦٠ مللي تسلا لمدة ١٢ ساعة اعلى قيم للصفات في كلا الحشرتين ادي التفاعل بين المجال المغناطيسي و الاصناف المختبره الى تحسين جميع الصفات مقارنة بالكنترول في كلا الحشرتين اظهرت اختلافات واضحة في صفات طول البادرات و الوزن الجاف للبادرات و دليل قوة البادرات و التوصيل الكهربائي في الاصناف الثلاثة المصابة بكلا الحشرتين. و قد اعطي مصر ١ اعلى القيم في طول البادره و دليل قوة البادرات و الوزن الجاف للبادرات يليه سدس ١٢ و سخا ٩٣ مع الحشرة ناقبة الحبوب الصغرى. بينما مع حشره سوسه الارز كان الترتيب سخا ٩٣ ثم سدس ١٢ و مصر ١ اعلى الترتيب . ادي تعريض البذور المصابة للمجال المغناطيسي لتأثير معنوي واضح في جميع الصفات مقارنة بالكنترول ظهر تأثير معنوي واضح نتيجة التفاعل بين المجال المغناطيسي و الاصناف على الوزن الجاف للبادرات و دليل قوة البادرات مع حشره سوسه الارز و ثاقبه الحبوب الصغرى بينما لا يوجد تأثير معنوي في نسبة الانبات و طول البادرات مع كلا الحشرتين و عموما جاء سخا ٩٣ في المركز الاول ثم مصر ١ و سدس ١٢ في انتاج اقل البادرات وزنا و اعلى قيم في دليل قوة البادرات تحت المعاملات المختلفة. اظهر التوصيل الكهربائي تأثير معنوي عالي باستخدام المجال المغناطيسي و كانت اقل القيم مع ٦٠ مللي تسلا لمدة ١٢ ساعة.صفات جوده البذور: - جوده البذور تشمل الرطوبة % , البروتين % , الدهون % , الرماد % و الكربوهيدرات % - اعطى استخدام المجال المغناطيسي ٦٠ مللي تسلا لمدة ١٢ ساعة زيادة طفيفة في الرطوبة % و الكربوهيدرات الكليه و نقص طفيف في البروتين % و الدهون % و الالياف %.