

## Sensitivity Injury *Tetranychus urticae* for Two Types of Sweet Pepper and the Effect of Infection on the Physical and Chemical Characteristics of the Plants.

Abd El- Rahman, H. A. and M. S. El-keblawy

Plant Protection Research Institute, Agricultural Research Center, Giza, Egypt.



### ABSTRACT

An experiment was conducted during two successive growing seasons of 2015 and 2016 under greenhouse condition ( four screen-houses 9m width x 40m length x 2m height) were established for investigate the susceptibility of two sweet pepper hybrids to infestation by two spotted spider mite *Tetranychus urticae*.The sweet pepper hybrids in) Kafr El-Sheikh-El-Hamol-El-Manwfa(showed different susceptibility degree to the aforementioned pest. The data showed that liebergen was the highest density population of *T.urticae* in seasons of 2015 and 2016 under greenhouse while steric was low density population of *T.urticae* .Liebergen was the highest sensitivity while steric was the least sensitivity.Out of the two hybrids only one steric rated as low resistant against the two spotted spider mites *Tetranychus urticae*. The remaining hybrids showed variable degree of resistance to two spotted spider mite *Tetranychus urticae*.Therefore this hybrids could be recommended for integrated pest management program .

### INTRODUCTION

Tetranychid mites are common pests in agricultural systems causing in many cases, greater economic losses than any other arthropod pests. The two-spotted spider mites, *Tetranychus urticae* is considered as one of the major pests attacking different agricultural crops such as field crops, vegetables, fruits and ornamental plants. The two-spotted spider mites *T. urticae* (Koch) has been extensively studied and the early work was reviewed by Huffaker *et al.* (1969). *T. urticae* infests a wide range of economic plants in the field such as cotton (Leigh *et al.*, 1968), strawberry (Sances *et al.*, 1981 and 1982), cucumber (De-ponti, 1980), tomato (Rodriguez *et al.*, 1972), peanuts (Boykin and Campbell, 1982), peppermint (Hollingsworth, 1980).

Sweet pepper is an important agricultural crop. Not only because of its economic importance, but also for the nutritional values of its fruits, mainly due to the fact that they are an excellent source of natural colors and antioxidant compounds (Howard *et al.*, 2000) In this respect, sweet pepper is considered an excellent source of bioactive nutrients, such as carotenoids, vitamin C and phenolic compounds(Navarro *et al.*, 2006).Hundreds varieties of peppers are now available for the greenhouses. They range widely in size, shape, color, flavor, disease resistance and season of maturity. However, many previous workers indicated that varieties of pepper plants play a great role for improving the growth and productivity (Rembalkowska *et al* 2005).

It is generally assumed that environmental factors and agricultural techniques have an effect on the quality of vegetables and fruits (Wang.,2006; Bafeel, and Ibrahim, 2008 and Nunez *et al* ,2011).In addition pepper is one of the most important exportable crops in Egypt. Pepper crops grown in greenhouses can be seriously damaged by various insects and mites. Salehi *et.al.*(2007) Sweet pepper have cytokinins and trace elements (Fe, Cu, Zn, Co, Mo, Mn and Ni) as well as vitamins and amino acids (Zodape *et al.* 2011).

This study aims to reduce the cost of populations of *T. urticae* and to allow the start of the sweet pepper growing season earlier under greenhouses with the two hybrids of sweet pepper plants considering growth.

### MATERIALS AND METHODS

#### Culture techniques:

The two-spotted spider mite, *Tetranychus urticae* (Koch) (Acarina: Tetranychidae),was reared according to Dittrich (1962) in colonies obtained from sweet pepper plants from greenhouse plants

#### Egg deposition and egg- hatching of *T. urticae*:

Five adult females' mites *T.urticae* of known age were placed on each disc. Numbers of eggs laid were assessed individually on different discs after 24 and 48 hours later Keratum *et al.* (1994). The number of hatched eggs was also counted 4 days after egg deposition. Assessment was conducted at 25± 2°C and 70±5 R.H. Each treatment was replicated four times. The same technique was repeated by sweet pepper leaf discs.

#### Antioxidant enzymes and chlorophyll (A and B) concentration

For enzyme assays in plants, 0.5 g leaf material was homogenized at 0-4°C in 3 ml of 50 mM TRIS buffer (pH 7.8), containing 1 mM EDTA-Na<sub>2</sub> and 7.5% polyvinylpyrrolidone. The homogenates were centrifuged (12,000 rpm, 20 min, 4°C), and the total soluble enzyme activities were measured spectrophotometrically in the supernatant (Hafez, 2010). All measurements were carried out at 25°C, using the model UV-160A spectrophotometer (Shimadzu, Japan). The enzyme assays were tested three times.

Activity of catalase (CAT) was determined spectrophotometrically according to Aebi (1984). Polyphenol oxidase (PPO) activity was determined according to the method described by Malik and Singh (1980). Changes in the absorbance at 495 nm were recorded every 30 sec intervals for 3 min. Enzyme activity was expressed as the increase in absorbance min<sup>-1</sup> g<sup>-1</sup> fresh weight. Peroxidase (POX) activity was directly determined of the crude enzyme extract according to a typical procedure proposed by Hammerschmidt *et al.*, (1982). Changes in absorbance at 470 nm were recorded every 30 sec intervals for 3min. Enzyme activity was expressed as the increase in absorbance min<sup>-1</sup> g<sup>-1</sup> fresh weight.

Chlorophyll (Chl.) concentration as mg/g fresh weight of one gram fresh leaves was extracted with 5 ml N,N-dimethyl-formamid for overnight at 5°C then estimated Chl. a and b spectrophotometrically at 663 and 647 nm as described by Moran and Porath (1982).

**Methods of determination NPK:**

Samples of leaves of (steric and liebergen) tissues were dried till constant weight. The dried leaves were grounded to fine powder.0.2gm of the fine was digested using sulphuric acid and perchloric acid (5:1) then the solution was completed to 50 ml using distilled water. The final solution was used to determine nitrogen, phosphorus and potassium as follows:

The nitrogen content in steric and liebergen leaves was determined in all of the before mentioned treatment by kjeldahl method according to Chalmers (1984).Phosphorus was determined according to Chapman and Pratt (1961).Potassium was determined using a flame photometer according to Chapman and Pratt (1961) Potassium percentage was determined by a flame photometer with aid of standard curve of known concentrations of potassium.

**Field experiment**

An experiment was conducted in a greenhouse at El-Manwfa village, El- Hamol district,Kafr El-Sheikh-governorate; during two successive growing seasons of 2015 and 2016 using new two hybrids sweet pepper for studying the population dynamics in hybrid, sweet pepper. The greenhouse was 510 m2 with 60m long and 8.5m wide,divided into 5 rides.Each ridge 90 cm.wide and 60m long. Seedling were transplanted on both sides of the ridges at 60 cm. apart nearly. A complete randomized block sides with three replicated for each hybrids were adopted, the tested hybrids which were distributed at random within the replicate. Each plot contained 10 plants thus the area of each ridge was 16.2 m2.All

received the stander cultivation practices that area including organic and mineral fertilization drip irrigation.

Data were statistically analyzed by using F test, means were compared according Duncan's multiple range test as described by Steel and Torrie (1982)and (Duncan,1955 )

Two spotted spider mite *T.urticae* populations was estimated on sweet pepper leaves of two hybrids. The numbers of *Tetranychus urticae* was weekly counted on the terminal leaf one each plant. Ten leaves were randomly selected from each plot and examinal using a hand lens.

**RESULTS AND DISCUSSION**

**1. Biology of the two-spotted spider mites *T.urticae*:-**

One of the important bases of the integrated pest management is the studying of the different effects of biological agents on mite egg deposition and the different responses of the eggs to these agents. These informations are of great importance for the entomologist to reach the different relations of certain importance to the phytophagous mites.

**2. Egg deposition of adult females of *T. urticae* on hybrids of sweet pepper:**

The data shown in Tables (1 and 2) indicated that the mean number of eggs deposited by adult female mites *T.urticae* on hybrids sweet pepper. The result in suggested that liebergen caused the highest reduction in egg deposition through the first day to five day (30.25: 95.50) followed by steric (20.75: 86.50 ). In general the effect of *T.urticae* can be arranged as steric > liebergen to egg deposition of adult females of *T. urticae* on hybrids of sweet pepper.

**Table 1. Egg deposition of adult females of *T. urticae* on hybrids of sweet pepper:**

Compounds	No. of egg deposited/5 adults					Mean±SD
	1 <sup>st</sup> day	2 <sup>nd</sup> day	3 <sup>rd</sup> day	4 <sup>th</sup> day	5 <sup>th</sup> day	
steric	20.75±0.95c	42.50 ±0.57c	57.00±81c	59.25±0.95c	86.50±1.29c	89.60±0.91c
Liebergen	30.25±1.70b	55.50±2.38a	67.25±2.06a	89.75±2.75a	95.50 ±1.29a	97.45±2.036a

**Table 2. Reduction of adult females of *T. urticae* on hybrids of sweet pepper:**

Compounds	Reduction%					Mean
	1 <sup>st</sup> day	2 <sup>nd</sup> day	3 <sup>rd</sup> day	4 <sup>th</sup> day	5 <sup>th</sup> day	
steric	71	62	48	21	19	44.2
Liebergen	88	72	60	38	31	57.8

**3. Population of adult mites *T.urticae* on sweet pepper during 2015 and 2016 seasons.**

The data were shown in Table (3) that liebergen was the highest density population of *T.urticae* in seasons of 2015 and 2016 under greenhouse while steric was low density population of *T.urticae* in seasons of 2015 and 2016.This experiment was carried out to sensitivity injury on mite populations of *T.urticae* so was the highest sensitivity while steric was the least sensitivity to the two hybrids of pepper.

On the other hand steric hybrid showed low liebergen number of *T.urticae* from month (march to October),the highest density population of *T.urticae* for hybrid steric(258) was observed in dune .While liebergen was the highest number of *T.urticae* from

month (march to October) while month June was the highest density population of *T.urticae* for hybrid liebergen (332) . Steric appeared high resistance but liebergen was the lowest resistance to *T.urticae* on hybrids sweet pepper Thus the author suggest the inclusion of this resistant sweet pepper hybrid in the integrated pest management program.

**Table 3. Population of adult mites *T.urticae* on sweet pepper during 2015 and 2016 seasons.**

Months	Mean No. <i>T.urticae</i> Season 2015		Mean No. <i>T.urticae</i> Season 2016	
	Mean No. <i>T.urticae</i> to steric	Mean No. <i>T.urticae</i> To Liebergen	Mean No. <i>T.urticae</i> to steric	Mean No. <i>T.urticae</i> To Liebergen
	March	6.0	8.0	4.0
April	8.0	17.0	11.0	18.0
May	156.0	233.0	188.0	212.0
June	258.0	332.0	276.0	307.0
July	194.0	212.0	183.0	203.0
August	125.0	163.0	107.0	112.0
September	36.0	54.0	33.0	46.0
October	32.0	44.0	15.0	33.0

**4. Determination of NPK in sweet pepper:**

Some investigators found that the supply and absorption of minerals were always positively correlated with pest infestation and that analysis of the host leaves may be valuable in nutrition studies of phytophagous pests (Highland and Roberts 1984 and Roinien and Tahvanaian 1989).

Thus the determination of such minerals as nitrogen, potassium and phosphorus in different host plant leaves used to rear. The data in Table (4) illustrate the rate of NPK for two host plants steric and liebergen. It is clear that steric has the highest rates of the three elements under study (6.888, 0.675 and 2.464) nitrogen, potassium and phosphorus. While liebergen plants have the lowest rates in the three elements (5.938, 0.464 and 1.833). The rate of potassium meq/ 100gm dry weight in the two host plants was steric and liebergen. The rate of nitrogen as NO<sub>3</sub> meq/ 100gm.d.w. in the two hybrids sweet pepper. The rate of phosphorus as meq/ 100gm.d.w. in the two hybrids sweet pepper was steric and liebergen.

**Table 4. Determination of NPK for steric and liebergen leaves.**

NO	N Mg/L	P Mg/L	K Mg/L
steric	6.888	0.675	2.464
Liebergen	5.938	0.464	1.833

The following points can be concluded from the results on the effect of different rates of the three elements in the two hybrids sweet pepper was steric and liebergen to spider mites *T.urticae*.

**5. Determination of antioxidant enzymes and chlorophyll (A and B) to sweet pepper hybrids.**

The data in figure (1 and 2) showed that estimation of enzymes in plants illustrated the extent of the difference between the different types and the extent of its relationship with the numbers of the population of the spider mite, steric higher than the kind of liebergen in the concentrations of enzyme and that meaning that the concentrations of enzyme is the reasons for the low number of population on the plants sweet pepper as Liebergen (yellow) the largest population and the ratio of the concentrations of enzymes and at least this shows that these concentrations have an inverse relationship existing population numbers. The results in figure (3) showed that steric was much higher degree contains phenols.

The data in figure (4) showed that the chlorophyll outside the scope of the competition between the two species in the presence of the population on sweet pepper. Park,(2002) chlorophyll content was reduced by approximately 55 and 80%, and greenness was reduced by approximately 50 and 80% by feeding by immature and adult *T. urticae*, respectively. Jack,(1983) chlorophyll was reduced in proportion to the degree of injury sustained.

**Fig. 1. Determination of antioxidant enzymes to sweet pepper hybrids.**

**Fig. 2. Determination of peroxidase activity to sweet pepper hybrids.**

**Fig. 3. Determination of polyphenol oxidase to sweet pepper hybrids.**

while liebergen was high resistance (HS) hybrids on basis of results.

**Table 5. Susceptibility degrees of *T.urticae* to the tested pepper hybrids during 2015 and 2016 seasons.**

NO	Hybrids	<i>Tetranychus.urticae</i>			Susceptibility degree
		2015	2016	Mean±SD	
1	Liebergen	4.24	4.48	4.28 ± 0.04	HS
2	steric	2.14	2.27	2.54 ± 0.42	S
-----	Mean	3.19	3.37	1.88	-----

HS – Highly susceptible                      S- Susceptible

It could be concluded that out of the two sweet pepper hybrids evaluated for their resistance hybrid was rated as resistant to *Tetranychus.urticae*. Thus the author suggest the inclusion of this resistant sweet pepper hybrid in the integrated pest management program.

**Fig. 4. Determination of chlorophyll (A) to sweet pepper hybrids.**

The data in figure(°) showed that steric of sweet pepper was higher than Liebergen and this shows that the chlorophyll (b) has to do with the number of inhabitants of the spider mite .Landeros(°) The data indicate that increased mite density coincides with a decrease in the net photosynthetic rate, transpiration and chlorophyll content. Higher mite densities on leaves cause stomata to remain open for longer periods, which allows a greater loss of water. Spider mite densities of 10 and 50 mites per leaf cause a reduction in flower stem length of 17 and 26%, respectively, as compared to plants with no mites present.

**Fig. 5. Determination of chlorophyll (B) to sweet pepper hybrids.**

**6. Population trend and susceptibility degrees of *T.urticae* to the tested pepper hybrids during 2015 and 2016 seasons.**

Out of the two sweet pepper hybrids liebergen hybrid had significantly high mean number (4.28) of spider mite per leaf as compared to the other tested hybrids steric with 2.54 spider mite per leaf ranked second in its susceptibility. Steric low numbers the pest (2.14 and 2.27 ) spider mite per leaf, respectively and appeared low resistance (S) hybrids on basis of results

**REFERENCES**

Aebi ,H.( 1984 ).(Catalase in vitro. Methods Enzymol . 127-131 :100

Bafeel, S.O. and M.M. Ibrahim, (2008). Antioxidants and accumulation of  $\alpha$ -tocopherol induce chilling tolerance in *Medicago sativa*. *Int. J. Agric. Biol.*, 10: 593–598.

Boykin, L. S. and W. V. Campbell (1982). Rate of population increase of the two-spotted spider mite (Acari: Tetranychidae) on peanut leaves treated with pesticides. *J. Econ. Entomol.* (75): 966-971.

Chalmers, R.A. (1984). Methods of proline analysis Publ.Chichester, Halsted press. Addivision of john Wiley Sons. New York, 371p.

Chapman, H.D. and P.F. Pratt (1961). Methods of analysis for soils, plants and waters. California. Univ. Devison of Agric. Sciences.

De-Ponti, O.M.B (1980). Resistance in *Cucumis sativus* L. to *Tetranychus urticae* Koch 6. Comparison of near isogenic bitter and non- bitter varieties for resistance. *Euphytica*, (29): 261-265.

Dittrich, V. (1962). A comparative study of toxicological test methods on a population of the two-spotted spider mite (*T.urticae*). *J. Econ. Entomol.* 55 (5) 644- 648.

Duncan, D.B (1955).; Multiple range and Multiple F tests. *Biometrics* 11:1-42.

Hafez, Y. M. (2010). Control of Botrytis cinerea by the resistance inducers benzothiadiazole (BTH) and hydrogen peroxide on white pepper fruits under postharvest storage. *Acta. Phytopathol. Entomol. Hung.*, 45 (1): 13-29.

Hammerschmidt, R., E.M. Nuckles and J. Kuć, ( 1982). Association of enhanced peroxidase activity with induced systemic resistance of cucumber to *Colletotrichum lagenarium*. *Physiol. Plant Pathol.*, 20(1):73-82.

Highland, A. and Robrts, U. (1984). Spider mite webbing II. The effect of webbing removal on egg hatchability. *Comparative Biochem Physiol* 51: 457- 462.

- Hollingsworth, C.S. (1980). Two-spotted spider mite, *Tetranychus urticae* in Oregon pepper mint, *Mentha piperita*: Sampling population dynamic and economic injury. Ph.D. Thesis, Oregon State University. Corvallis, pp. 159.
- Howard, L.R., S.T. Talcott, C.H. Brenes and B. Villalon. (2000). Changes in phytochemical and antioxidant activity of selected pepper cultivars (*Capsicum* species) as influenced by maturity. *J. Agric. Food Chem* 48:1713-1720.
- Huffaker, C. B.; M. Van De Vrie and J.A. Me Murtry (1969). The ecology of tetranychid mites and their natural control. *A Rev. Entomol*,(14): 125-174.
- Jack, A.F. (1983). Comparison of two Phytoseiulus species as predators of twospotted spider mites on greenhouse ornamentals. *Environmental Entomology* 10: 524-527.
- Keratum, A., Y.; H. Anber; M.M. Essawy and F.L El-Shahawi (1994). The effect of permethrin residues on the activity of the predators *Phytoseiulus persimilis* and *Amblyseius fallacis*. *Alex. Sci. Exch*, 15(1): 67-82.
- Landeros, A, C (2015). Biology and natural enemies of *Tetranychus urticae* in sweet pepper. *Manejo Integrado de Plagas y Agroecología. Centro Agronomico Tropical de Investigacion y Ensenanza, Turrialba, Costa Rica*, 74: 34-40.
- Leigh, T.F.; R.E. Hunter and A.H. Hyer (1968). Spider mite effects on yield and quality of four cotton varieties. *Calif. Agric.* (22): 4-5.
- Malik, C. P. and M. B. Singh. (1980). In: *Plant Emynology and Histoenzymology*. Kalyani Publishers. Indian and printed in Navin. Shanndara. Delhi PP. 54-56.
- Moran, R. and D. Porath, (1982). Chlorophyll determination in intact tissues using N,N-Dimethyl formamide. *Plant Physiol.*, 69: 1370-1381.
- Navarro, H.;J. Demi, S. Uzun and V. Ceyhan, (2006) The effects of different greenhouse covering materials on energy requirement, growth and yield of aubergine. *Energy*, Vo. 31 (12): 178-1788.
- Nunez, D; Daniel; J. Grimaldo and C.D. Lourdes (2011). Nitrogen fertilization effect on Antioxidants Compounds in Fruits of Chili Pepper *Capsicum chinense*/ *Int. J. Agric. Biol.*, Vol. 13.
- Park, J.G (2002). Large scale production and release of the predatory mites *Pytoseiulus macropilis* (Banks) to control *Tetranychus urticae* Koch on commercial strawberry plantations. *Bull. Ent. Soc. Egypt, Econ. Ser. Num.* 33.153-163.
- Rembińska E., Hallmann E., Szafirowska A. (2005). Nutritive quality of tomato fruits from organic and conventional cultivation. *Culinary Arts & Sciences V. Warsaw Agric. Univ. Poland*: 193-202.7.
- Rodriguez, J.C.; D. E. Knavel and O. J. Aina (1972). Studies in the resistance of tomato to mites. *J. Econ. Entomol*, (65): 50-53.
- Roinien, V.J and Tahvanainen, J.Y (1989). Protected cultivation of sweet pepper hybrids under net-house in Indian conditions. *Acta Horticulturae*, 659: 515-521.
- Salehi, M., K. Izadpanah and M. Siampour, (2007). Characterization of a phytoplasma associated with cabbage yellows in Iran. *Plant Dis.*, 91:625-630.
- Sances, F.V.; J.A. Wyman; LP. Ting; R. Van Steenwyk and E.R. Oatman (1981). Spider mite interactions with photosynthesis, transpiration and productivity of strawberry. *Environ. Entomol*, (10): 442-448.
- Sances, F.V.; N.C. Toscano; E.R. Oatman; L.F. Lapre; M.V. Johnson and V. Voth (1982). Reduction in plants processes by *Tetranychus urticae* feeding on strawberry. *Environ. Entomol.*(11):733-737.
- Steel, Y.F. and Torrie, H. N. (1982). Protected cultivation of sweet pepper hybrids under net-house in Indian conditions. *Acta Horticult.* 659: 515-521
- Wang, S.Y., (2006). Effect of pre-harvest conditions on antioxidant capacity in fruits. *Acta Hort.*, 712: 299-305.
- Zodape, D.; Y. Sandeep K., Dhillon, T. S and Parminder, S (2011) Protected cultivation of sweet pepper hybrids under net-house in Indian conditions. *Acta Horticult.* 659: 515-521.

## حساسية الإصابة بالعنكبوت الأحمر العادي لصنفين من الفلفل الحلو وتأثير الإصابة على الصفات الطبيعية والكيميائية للنباتات.

حمدي عبد الرحيم عبد الرحمن و محمود سعد القبلاوي  
معهد بحوث وقاية النباتات - مركز البحوث الزراعية - وزارة الزراعة

أجريت التجربة خلال موسمي ٢٠١٥ و ٢٠١٦ تحت الصوب الزراعية في قرية المناوفة مركز الحامول محافظة كفر الشيخ ومعهد وقاية النباتات فرع سخا (مركز البحوث الزراعية) لدراسة حساسية صنفين من الفلفل الحلو وهما (لامبرجيني و استيرك) بالإصابة بالعنكبوت الأحمر العادي وأظهر الصنف استيرك حساسية عالية بالمقارنة بالصنف لامبرجيني تجاه الأكاروس الأحمر حيث اتضح أنه منخفض المقاومة وكان الصنف استيرك أعلى مقاومة في التجربة محل الدراسة حيث كانت الكثافة الأكاروسية للصنف لامبرجيني أعلى بالمقارنة بالصنف استيرك في الموسمين وكانت مقاومة الصنف استيرك للأكاروسات ناتجة من الاختلافات في الصفات الطبيعية والكيميائية بين الصنفين لنبات الفلفل الحلو محل الدراسة وذلك ينصح بزراعة الصنف المقاوم واستخدامه في برامج المعالجة المتكاملة للأكاروسات تحت البيوت المحمي