

PHYSIOLOGICAL AND ULTRASTRUCTURE RESPONSES  
OF *Schistocerca gregaria* (FORSKAL) TO *Metarhizium*  
*acridum* AND FOUR BIO ACTIVE COMPOUNDS

Farrag, A. A.<sup>1</sup>; T. A. Abd-Elfattah<sup>2</sup>; G. M. Abdelatef<sup>2</sup>  
and

M. Kh. El-Dydamony<sup>2\*</sup>

1-El Azhar University, Faculty of science, Microbiology Dep.  
Madent Nasr, Cairo

2-Plant Protection Research Institute, locust and  
Grasshopper Research Dep. Doki, Giza.

\*khyri\_dedo@yahoo.com



## ABSTRACT

The effect of *Metarhizium acridum* and four bioactive compounds (Neem, L- Glutamic acid, *Schinus molle* and abamectin) in sole treatments and in combination with *M. acridum* was studied on the mortality of 5<sup>th</sup> nymphal instar of desert locust *Schistocerca gregaria* as well as the time mortality responses and the ultra-structure of females ovary. The obtained results showed that mortality in case of *Metarhizium acridum* treatment reached to 100% after 14 days, while in case of neem treatment the mortality reached to 60 % after 12 days, at the same time the mortality reached up to 20 % after 10 days post treatment with L- Glutamic acid, but such mortality reached to 50 % after 14 days post treatment with *Schinus molle* extract, finally the mortality reached to 60 % after 11 days of abamectin treatment. All the mixtures with *M. acridum* caused 100 % mortality by the 5<sup>th</sup> day post treatment. However the sole treatments caused long time to kill 50 % of treated insects (LT<sub>50</sub>), where the LT<sub>50</sub> of *M. acridum* was 7.261 day as the fastest treatment, while all mixtures caused accelerate in the mortality, but the best result was obtained in case of *M. acridum* + neem treatment which LT<sub>50</sub> was only 1.625 day.

Any way there were different effects on ultra-structure of the ovary as result of treatments used. Such effects varied between changes in nucleus shape to changes and destruction of nucleus itself.

**Keywords:** *Schistocerca gregaria*, *Metarhizium acridum*, plant extract, amino acid, L- Glutamic acid, *Schinus molle*, abamectin, bioassay, time mortality responses and ovary ultrastructure.

## INTRODUCTION

The desert locust *S. gregaria* has been the most serious crop pest in many countries of Africa and Asia, it is major pest of many tropical and subtropical countries causing extensive damage to the foliar part of many plants particularly during years with locust outbreaks. (Abd El-Fattah, 2005 and Ceccato, *et al.*, 2007). The current method for locust control based on applying synthetic chemical insecticides (Gamal *et al.*, 2012). Entomopathogenic fungi *Metarhizium acridum*, has been commercialized and used successfully for biocontrol of grasshopper pests in Africa and Australia (Stuart *et al.*, 2012) now produced commercially with trade names such as Green Muscle and Green Guard), characterized as safe pathogen to non-target organisms, and they can penetrate directly through the cuticle and

not necessarily need to be ingested in order to initiate disease (Goettel et al., 1990) The application of Green Muscle considered as a new technique to locusts and grasshoppers control (Mahgoub et al., 2011). Neem tree *Azadirachta indica* has been used for many years against several insect pests (Hamadah et al., 2013). Neem tree has strong insecticidal activity against *S. gregaria* (Nicol and Schmutterer, 1991). *S. molle* has active substances, such as terpenes, tannins, alkaloids, flavonoids, saponins, gums, linoleic acid, oleoresins, mostly in leaves and fruits (Hayouni et al., 2008), Abamectin is a natural product of the soil microorganism *Streptomyces avermitilis* and shows spectrum insecticide and acaricide with high pesticidal activity, (Qiao et al. 2012). Abamectin and enhanced the effect of *M. acridum* against desert locust nymphs in the field (Mohamed et al. 2014). L-Glutamic acid is known as excitatory transmitter at the neuromuscular junction of invertebrates (Wafford and Sattelle, 1989), while (Clements and May, 1974) showed that when *Schistocerca gregaria* nerve-muscle exposed to glutamate caused a variety of responses, some of which were shown to be abnormal and were much more severely affected. *S. gregaria* had panoistic ovary, which is composed of number of ovarioles (Martoja, 1964). Each ovariole was surrounded by basal lamina and consists of follicle cells and oocytes. The aim of present work is study the effect of four bioactive compounds (at rate of 1 tenth of recommended dose of each compound) on acceleration mortality caused by *M. acridum* to achieve safe desert locust control procedures, as well as the effect of such treatments on the ultra-structure of female ovarian tissues.

## MATERIALS AND METHODS

### Desert locust nymphs

Fifth nymphal instar of *S. gregaria* two days after final molting, were kindly obtained from desert locust colony maintained in Locust and Grasshoppers Research Dep., Plant Protection Research Institute, ARC, Cairo Egypt. Desert locust individuals were reared in the laboratory according to (Robert et al., 2002). The colony was fortified with wild insect collect from the field each year.

### *Metarhizium acridum*

The entomopathogenic fungus used during the study is *M. acridum* (formerly *Metarhizium anisopliae* var. *acridum*) Bischoff et al., 2009, isolate (IMI330189) was kindly obtained from BASF, South Africa, under the commercial name Green Muscle®. The spores were suspended in sterilized water; trace of Tween (80) was added. The concentration was adjusted to  $5 \times 10^8$  spores/ml, each nymph received 5µl of the final solution.

### Neem (Azadirachtin)

Azadirachtin, under the commercial name Safe-oil 0.03 % EC, at concentration of 1ml/liter distilled water.

### *Schinus molle* extract

50 g of fresh aerial part of *S. molle*, were air-dried at lab temperature, dried in oven at 40°C till constant weight then ground to fine powder, add to

800 ml liter of distilled water in volumetric flask for 3 days and repeated 4 times, then filtrated. Combined filtrates were evaporated under reduced pressure using rotary evaporator apparatus until a minimum amount of solvent remained which gives (3 g) at last. The extract (brownish sticky) was stored in a refrigerator at 5 °C and kept for using in different analysis. The concentration used was 1 g from extract added to 100 ml distilled water (Woo *et al.*, 1977).

#### **Amino acid (L-Glutamic acid)**

Molare solution from L-glutamic acid is prepared by addition of 147.13 gm \ litter distilled water. concentration are used as 0.1 from molar solution (Krasilnikov and Bakhrarov, 1983) each nymph received 5µl of the final solution.

#### **Abamectin**

5-O-demethylavermectin A<sub>1a</sub> (i) mixture with 5-O-demethyl-25-de(1-methylpropyl)-25-(1-methylethyl) avermectin A<sub>1a</sub> (ii). Under the commercial name, Agromic 1.8 % EC, was used at concentration of 1ml/liter distilled water, each nymph received 5µl of the final solution.

#### **Mixtures**

*Metarhizium acridum* was used in combination with Neem, *S. molle*, L-Glutamic acid and Abamectin, the mixture solution contain the same concentration for each compound as in sole case.

#### **Treatments**

Thirty individuals of 5th instar nymphs were used in each treatment, divided into 3 replicates each. *M. acridum*, L-glutamic acid and Abamectin treatments were used as topical application, while Neem and *S. molle* were used as follow: 40 g of fresh clover were dipped in 100 ml of each used concentration, dried for 1 hr. in room temperature, then introduced to the nymphs. The mixtures were used as topical application, but in case of Neem and *S. molle* the nymphs were treated with *M. acridum* first then were feed on treated clover as described previously. Treated and untreated nymphs were kept in cages under laboratory conditions at 31 ± 0.5°C, also were feed and cleaned daily, mortality was observed and recorded daily. Extra five female 5<sup>th</sup> nymphal instar per each treatment were treated as described previously, then at the 4<sup>th</sup> day post treatment, these nymphs were subjected to dissection for Electron Microscope examination.

#### **Transmission Electron Microscopy (TEM) examination**

The insects were killed by twisting the head to break the "neck" membrane. The posterior tip of the abdomen was cut off and the head, with the gut attached, was removed. Ovary cleaned from the surrounding fat body and then it was dissected in ice-cold (0-5 °C) Karnovsky fixative, pH 7.3 (Karnovsky, 1965). The tissue pieces were washed twice in a buffer for 30 min. The specimens were then dehydrated in grades of ethanol; 50, 70, 80, 90 and 100%. The specimens were cleared in toluene for 10 min and then embedded in the resin of choice Epon. Semithin sections are cut from these blocks (stained with toluidine blue) and examined by the light microscope (Spurr, 1969). Ultrathin sections obtained from selected blocks were mounted on copper grids stained with uranyl acetate and lead citrate and then examined with Jol 1010 transmission electron

microscope (Reynolds, 1963). This technique was carried out at Regional Center for Mycology and Biotechnology, Al-Azhar University, Madenit Nasr, Cairo.

#### Statistical analyses

Mortality data were subjected to probit analyses according to Finney (1971) to calculate time mortality responses and its regression lines.

## RESULTS

Fig. (1) shows the accumulative mortality caused by *Metarhizium acridum* Neem, *Schinus molle*, L- Glutamic acid and abamectin used separately or in combination against *Schistocerca gregaria* 5<sup>th</sup> nymphal instar. It's clear that *M. anisopliae* treatment caused slow mortality progress began by day 5, ended with 100 % of mortality by day 14 post treatment, while other treatments (used at 1 tenth of the recommended dose) showed same trend, where mortality in Neem treatment began by day 4 ended with 60% mortality by day 12, while *S. molle* mortality initiated by day 6 post treatment ended with 50 % by day 14, also in case of L- Glutamic acid the mortality began by day 6 ended with 20 % by day 10. Finally in abamectin treatment the mortality began little bet early by day 3 post treatment ended with 60 % by day 11.

While all mixtures seriously increased the efficacy of *M. anisopliae*, the mortality began by the first day post treatment, the mortality % were (10, 22, 28 and 24%) respectively for the mixture of *M. anisopliae* with *Schinus molle*, L- Glutamic acid, abamectin and Neem, respectively. Such mortality ended by day 5 with 100 %.

In the same line with these data fig. (2) and table (1) demonstrate mortality time responses of the previous treatments, it could be concluded that sole treatments: *M. anisopliae*, Neem, *Schinus molle*, L- Glutamic acid and abamectin needed long period to kill half of the population ( $LT_{50}$ ). These  $LT_{50}$ s were 7.261, 10.489, 14.608, 46.568 and 9.055 days respectively, while  $LT_{90}$ s of the same treatments were 12.613, 27.815, 39.972, 462.937 and 29.907 days, respectively.

Although, mixtures greatly accelerate the mortality where  $LT_{50}$ s were 1.625, 2.414, 1.963, and 1.888 days for the mixtures of *M. anisopliae* with Neem, *Schinus molle*, L- Glutamic acid and abamectin, respectively,  $LT_{90}$ s of same treatments were 5.818, 5.496, 4.749, and 5.715 days, respectively.




Fig. (1) Accumulative mortality due to *Metarhizium acridum* Neem, L-Glutamic acid, *Schinusmolle* and abamectin treatment that are used separately or in combination against *Schistocerca gregaria* 5<sup>th</sup> nymphal instar.




Fig. (2) Time mortality responses of *Metarhizium acridum* Neem, L-Glutamic acid, *Schinusmolle* and abamectin Treatment that are used separately or in combination against *Schistocerca gregaria* 5<sup>th</sup> nymphal instar.

**Table (1) Time mortality responses of *Metarhizium acridum* Neem, L- Glutamic acid, *Schinus molle* and abamectin treatment that are used separately or in combination against *Schistocerca gregaria* 5<sup>th</sup> nymphal instar.**

Line name	LT <sub>50</sub> (Days)	Index	Slope	LT <sub>25</sub> (Days)	LT <sub>90</sub> (Days)
<i>M. acridum</i>	7.261	22.38	5.344	5.43	12.613
Neem	10.489	15.492	3.026	6.278	27.815
<i>S.mlle</i>	14.608	11.124	2.932	8.6	39.972
Amino acid	46.568	3.49	1.285	13.904	462.937
abamectin	9.055	17.946	2.47	4.828	29.907
<i>M. acridum</i> + neem	1.625	100	2.314	0.831	5.818
<i>M. acridum</i> + <i>S.mlle</i>	2.414	67.316	3.587	1.566	5.496
<i>M. acridum</i> + Amino acid	1.963	82.781	3.34	1.233	4.749
<i>M. acridum</i> + apamectin	1.888	86.07	2.664	1.054	5.715

### Transmission Electron Microscopy (TEM) examination

Electron microscope of ovarioles for untreated insects illustrated in Fig. 3<sub>(1)</sub>, showed nucleus (N) which coated with nucleus membrane (Nm) and contain normal chromatin (Ch), cytoplasm have normal organelles as rough endoplasmic reticulum (RER), mitochondria (M), and Golgi bodies (Gb). Yolk bodies (Y) are appearing black.

While in case of treated insects with *M. anisopliae* illustrated in Fig. 3<sub>(2)</sub>, it's clear that treatment caused rupture in nucleus (RN), malformation in mitochondria, cytoplasm was damaged and vasculization (V) ocured. The fungus hypha were clearly visible and wiedy spread in the cytoplasm which cause subversion of ovarioles.

Fig. 3<sub>(3)</sub>, shows ovary of treated insects with neem it's obvious that the shape of nucleus was irregular, the majority of chromatins were disappeared except gathering of few number in two spot, yolk was destroyed and decreased in size into small particles, while cytoplasm was destroyed and become incomprehensible, mitochondria were absent, and vasculization was ocured.

Ovarioles of insects treated with L- Glutamic acid illustrated in Fig. 3<sub>(4)</sub>, showed nucleus destruction, with dark abnormal chromatin materials, the yolk was disappeared, while the mitochondria were enlarged, the cytoplasm was damaged and vasculization ocured near basement membrane (Bm).

Fig. 3<sub>(5)</sub>, Illustrates ovarioles of treated insects with *Schinus molle* extract it's clear that the yolk was destroyed while spreation of numerous numbers of lipid droplets (L) were obvious, and vasculization was ocured and the mitochondria were absent.

Fig. 3<sub>(6)</sub>,Shows the ovarioles of treated insects with abamectin, it reveal that there were varying in shape and size of nucleus, while in some nucleus the chromatin became abnormal, the yolk destroyed and decreased in size also the cytoplasm was destroyed, the mitochondria was varied in size and shape finally vasculization ocured with large size.

Electron micrograph of ovarioles of treated insects with mixture from *M. anisopliae* and neem are illustrated in Fig. 4<sub>(7)</sub>,showed destruction of nucleus membrane, while the yolk was decreased in size and became pastel,

cytoplasm was destroyed, vasculization occurred finally the hypha were spread in ovarioles.


Electron micrograph of ovarioles for insects treated with mixture from *M. anisopliae* and L- Glutamic acid illustrated in Fig. 4<sub>(8)</sub>, showed destruction of cytoplasm (DC) in the same time cytoplasm organelles disappeared, as well as the yolk, vasculization occurred and differed in size, the spread of hypha in ovarioles were obvious.

Fig. 4<sub>(9)</sub>, displays electron micrograph of ovarioles of treated insects with *M. anisopliae* and *S. molle* extract mixture it's clear that the yolk differed in size, vasculization occurred and the hypha were spread inside the ovarioles.

Ovarioles of treated insects with *M. anisopliae* and abamectin mixture demonstrated in Fig. 4<sub>(10)</sub>, showed destruction of the nucleus, although the chromatin congregated in a spot, while the cytoplasm was destroyed, also mitochondria was absent, but small vasculization occurred and hypha were appeared near the nucleus.


## DISCUSSION

Many studies were done to evaluate the ability of *M. anisopliae* var. *acridum* to integrate with its other control agents in order to increase efficacy of *M. anisopliae* var. *acridum* for desert locust control operations *i.e.* integration with some insect growth regulators (IGRs) and antifeedant (El-Gamal *et al.*, 2004), and with abamectin and D-limonene (Mohamed *et al.*, 2014). Fungi usually cause insect mortality by one or more of the following: nutritional deficiency, invasion and destruction of tissues, and release of toxins. Fungal species have numerous strains that differ in their virulence and pathogenicity. The pathogenicity of fungus may be associated with the production of enzymes and mycotoxins during the course of infection in an insect Tanada & Kaya (1993). abamectin showed slow act against desert locust, this may be due to its mode of action where Wolstenholme and Rogers 2005 indicated that, it is likely that abamectin bind to multiple sites (including glutamate and GABA) in insect chloride channels. In general, the chloride ion flux produced by the opening of the channel into neuronal cells results in loss of cell function and disruption of nerve impulses. Consequently, invertebrates are paralyzed irreversibly and stop feeding. *Schinus molle* have diverse properties, such as insecticidal and repellent effects in different insects (Ferrero *et al.*, 2006, 2007; Abdel-Sattar *et al.*, 2009). *M. anisopliae* strain is compatible with neem and reduces survival of mosquito *Anopheles gambiae* adults after spraying (Fawrou *et al.*, 2012).



**Fig.(3):**1-:6 Electron micrograph of ovarioles 1-normal 2- treated with *M. anisopliae* 3- treated with neem 4-treated with L- Glutamic acid 5-treated with *S. molle*.6- treated with abamectin.

basement membrane (**Bm**) chromatin (**Ch**) endoplasmic reticulum (**RER**)  
golgi bodies (**Gb**) hypha (**H**) lipid droplets (**L**) mitochondria (**M**)  
nucleus (**N**) nucleus membrane (**Nm**) rupture in nucleus (**RN**)  
vasculization (**V**)



**Fig.(4):** 7-10 Electron micrograph of ovaries **7**-treated with mixture of *M. anisopliae* and neem **8**- treated with mixture of *M. anisopliae* and L-Glutamic acid. **9**- treated with mixture of *M. anisopliae* and *S. molle*. **10**-treated with with mixture of *M. anisopliae* and abamectin. basement membrane (**Bm**) destroyed cytoplasm (**DC**) chromatin (**Ch**) endoplasmic reticulum (**RER**) golgi apparatus (**Ga**) hypha (**H**) lipid droplets (**L**) mitochondria (**M**) nucleus (**N**) nucleus membrane (**Nm**) rupture in nucleus (**RN**) vasculization (**V**)

Mohamed, *et al.*, (2008) examined the ovarian follicle of adult *S. gregaria* by transmission electron microscopy and showed that, the follicular epithelium of vitellogenic follicles is composed a single layer of more or less columnar to cuboidal cells without nuclei. While Tobe and Pratt (1975) observed four distinct periods, an early growth period, previtellogenic period, vitellogenic period and finally chorionation, which precedes ovulation in *S. gregaria*. During vitellogenesis reserve materials were deposited in ooplasm and consequently oocytes become voluminous and filled with numerous yolk spheres (Simiczyjew and Margas 2001). John and James (1989) observed that the apical ooplasm of *S. gregaria* was rich in ribosomes and mitochondria, while rough endoplasmic reticulum (rER) and golgi complexes occupied the basal and basolateral regions of the cells. These ultra structures point to an increase in the activity of follicular cells associated with the synthesis and secretion of egg precursors. Similarly, newly emerged 5th instar nymphs of *S. gregaria* treated with cascade, rice bran extract and karate, synthetic pyrethroids each at LC<sub>50</sub> produced disturbance in protein synthesis of the ovary, which reflected an inhibition of ovarian maturation and showed a degeneration of ovarioles and oocytes, disintegrated mitochondria, enlarged vacuoles and cracked yolk bodies mostly in two halves (John and James., 1989), (Ferenz., 1993) and (Hussein *et al.*, 2008).

All treatments effect on ovarioles of *S. gregaria*, this effect vary from treatment to other as damaged in cytoplasm, uneven shape of nucleus, rupture in nucleus, destroyed in nucleus membrane, disappeared of chromatin, malformation in mitochondria, destroyed of yolk, lipid droplets, vasculization, and hypha which spread in ovarioles in treatment of *M. anisopliae* in alone case or in mixture with other treatments. All this unnatural phenomena may be due to effect of all treatments on central nerve cord by plugging of sodium and potassium channels, which ultimately stimulate the endocrine system and later on it may affect development of reproductive system, this result are in parallel to (Gupta *et al.*, 1992) and(Gupta, 1979).

## REFERENCES

- Abd El-Fattah, T. A. (2005). The combined effects of the entomopathogenic fungus *Metarhizium anisopliae* var *acridium* IMI 330189 (Green Muscle) and the sub-lethal doses of some insecticides on the desert locust, *Schistocerca gregaria* (Forsk.). Egypt. J. Agric Res., 83(2) 551-561.
- Abd El-Sattar, E.; Zaitoun, A.; Farag, M.; El-Gayed, S. and Harraz, F. (2009). Chemical composition, insecticidal and insect repellent activity of *Schinus molle* L. leaf and fruit essential oils against *Trogoderma granarium* and *Tribolium castaneum*. Nat. Prod. Res. 25, 1-10.
- Asi, M. R.; Bashir, M. H.; Afzal, M.; Ashfaq, M. and Sahi, S. T. (2010). Compatibility of entomopathogenic fungi, *Metarhizium anisopliae* and *Paecilomyces fumosoroseus* with selective insecticides. Pak. J. Bot, 42, 4207- 4214.

- Belhamel, K.; Abderrahim, A. and Ludwig, R. (2008). Chemical composition and antibacterial activity of the essential oil of *Schinus molle* L. grown in Algeria. *International Journal of Essential Oil Therapeutics* 2, 175-177.
- Bischoff J.F., Rehner S.A. and Humber R.A. (2009). A multilocus phylogeny of the *Metarhizium anisopliae* lineage. *Mycologia* 101 (4): 512–530
- Ceccato, P.; Cressman, K.; Giannini, A. and Trzaska, S. (2007). The desert locust upsurge in West Africa (2003–2005): Information on the desert locust early warning system and the prospects for seasonal climate forecasting. *Intl J Pest Management.*;53(1): 7-13.
- Clements, A. T. and May, T. E. (1974). Studies on locust neuromuscular physiology in relation to glutamic acid. *J. exp. Biol.* 60, 335–378.
- Engelmann, F. (1970). The physiology of insect reproduction. Department of zoology, Univ of California, Los Angeles, Engel. Pergamon: 307 pp.
- Fawrou, S.; Mady N.; Oumar F. and Jos'e M. A. (2012). Evaluation of Entomopathogenic Fungus *Metarhizium anisopliae* Formulated with Suneem (Neem Oil) against *Anopheles gambiae* s.l. and *Culex quinquefasciatus* Adults. *Malaria Chemotherapy, Control & Elimination*. Vol. (1), Article ID 235494, 6 pages.
- Fenske, R. A.; Black, K. G.; Elkner, K. P.; Lee, C. L.; Methner, M. M. and Soto R. (1990). Potential exposure and health risks of infants following indoor residential pesticide applications. *Am J Public Health.* ;80(6):689–693.
- Ferez H. J., (1993). Yolk protein accumulation in *Locusta migratoria* oocytes. *Journal of Insect Morphology and Embrology*, 22(2-4), 295-314.
- Ferrero, A.; Werdin, J. and Sánchez C. C. (2006). Actividad biológica de *Schinus molle* en *Triatoma infestans*. *Fitoterapia* 77, 381-383.
- Finney, D. J. (1971). *Probit Analysis*, Third Edition, London: Cambridge University Press.
- Gamal, A. H.; Salwa, S. R.; Eman, M. R. and Fatma H. (2012). Fungi Associated with the Desert Locust, *Schistocerca gregaria* (Forskål) and Their Laboratory Potency against Locust Nymphs. *Journal of Applied Sciences Research*, 8(8): 3914-3920.
- Goettel, M. S.; Poprawski, T. J.; Vandenberg, J. D. Li, Z., Roberts DW (1990). Safety to nontarget invertebrates of fungal biocontrol agents. In: "Safety of microbial insecticides", Laird M, Lacey LA, Davidson EW (eds.). Boca Raton, FL: CRC Press, pp. 209-231.
- Gupta H. C. L. (1999). *Insecticides: Toxicology and uses* (Agrotech publishing academy: Udaipur 92-99 pp.
- Gupta H. C. L., Singh R., Jain R. L. and Saxena R. C. (1992). Relative and residual toxicity of some synthetic pyrethroids, *Indian Journal of Entomology*, 54(1),. 34-38.
- Hamadah, Kh. Sh.; Ghoneim, K. S.; El-Hela, A. A.; Amer, S. M. and Mohammad, A. A. (2013). Disturbed Survival, Growth and Development of the Desert Locust *Schistocerca gregaria* by Different Extracts of *Azadirachta indica* (Meliaceae) and *Nigella sativa* (Ranunculaceae). *Egypt. Acad. J. Biolog. Sci.*, 6(2): 1 -21.

- Hayouni, E.; Chraief, I.; Abedrabba, M.; Bouix, M.; Leveau, J.; Mohammed, H. and Hamdi, M. (2008). Tunisian *Salvia officinalis* L. and *Schinus molle* L. essential oils: their chemical compositions and their preservative effects against *Salmonella* inoculated in minced beef meat. *Int. J. Food Microbiol.* 125, 242-251.
- Hussein M .A., Hamouda L. S., Bakr R. F.A., qH. A. and Elsokary Z. F (2008). Effect of Cascade, *Oryza sativa* bran extract and Karate on fine structure of the ovary of *Schistocerca gregaria*. *Egypt. Acad. J. Biol. Sci.* 1(1), , 13 – 21.
- John C. D. and James T. B. (1989). Ovarian follicle development during vitellogenesis in the house cricket, *Acheta domesticus*. *Journal of Morphology*, 200, 185-198.
- Karnovsky, M.J. (1965). The localization of cholinesterase activity in rat cardiac muscle by electron microscopy. *J. Cell. Biol.* 23, 217-232.
- Krasilnikov and A. Bakhramov (1983). Studies on the Effect of Glutamic, Aspartic and  $\gamma$ -Amino Butyric Acids on Locust Muscle Fibre Membrane Potential. *Gen. Physiol. Biophys.* (1983), 2, 133—135.
- Mahgoub M. M.; Hamadttu A. E.; and Magzoub, O. B. (2011). Use of teflubenzuron alone and combined with *Metarhizium anisopliae* and Phenylacetone nitrile as control agent against the desert locust, *schistocerca gregaria* (Forsk.) (*Orthoptera: acrididae*).
- Martoja, R. (1964). Untype particulier d'appareil genital femelle chez les insects: les ovarioles adenomorphes du coleopteran, *Steraspis speciosa*. *Bull. Soc. Zool. Fr.* 89: 614-41.
- Mohamed A. H.; Laila S. H.; Reda F. A. B.; Hassan H. A.; and Elsokary Z. F. (2008). Effect of Cascade, *Oriza sativa* bran extract and Karate on fine structure of the ovary of *Schistocerca gregaria*. *Egypt. Acad. J. biolog. Sci.*, 1 ( 1) 13 – 21.
- Mohamed, Gehan A., Hala M. Ibrahim and G. M. Abdelatef (2014). Novel pesticides for desert locust *Schistocercagregaria* (Forsk.) control. *Journal of Plant Protection and Pathology Mansoura Univ.* 5: 1065-1071.
- Nicol, C. M. Y. and Schmutterer, H. (1991). Contact effects of seed oil from the neem tree *Azadirachta indica* (A. Juss.), on nymphs of the gregarious phase of the desert locust, *Schistocerca gregaria* (Forsk.). *J. Appl. Entomol.*, 111 (2): 197-205.
- Qiao K., X. Liu, H. Wang, X. Xia, X. Ji and K. Wang (2012). Effect of Abamectin on root-knot nematodes and tomato yield. *Pest Management Science.* 68 (6): 853-857.
- Reynolds, S.E., (1963). The use of lead citrate at high pH as an electron opaque stain in electron microscopy. *J. Cell Biol.*, 1: 208
- Robert M. Ouedraogo, Andrena Kamp, Mark S. Goettel, Jacques Brodeur and Michael J. Bidochka (2002). Attenuation of fungal infection in thermoregulating locust migratoria is accompanied by changes in haemolymph proteins. *Journal of Invertebrate Pathology* 81: 19-24.
- Simiczjew B. and Margas W. (2001). Ovary structure in the bat flea, *Ischnopsyllus* spp. (Siphonaptera: Ischnopsyllidae), Phylogenetic Implications, *Zoolog. Polon.*, 46(1-4), 5-14.

- Spnrr, A. R., (1969). A low-viscosity epoxy resin embedding medium for electron microscopy. / *Ultrastructure Res.*, 26:31-43.
- Steinbauer, M. and Wanjura, W. (2002). Christmas beetles (Anoplognathus spp., Coleoptera: Scarabaeidae) mistake peppercorn trees for eucalypts. *J. Nat. Hist.* 36, 119-125.
- Stuart, B. Krasnoff, Ulrich English, Paula G. Miller, Michael L. Shuler, Raymond P. Glahn, Bruno G. G. Donzelli and Donna M. Gibson (2012). Metacridamides A and B, Macrocycles from Conidia of the Entomopathogenic Fungus *Metarhizium acridum*, *Journal of National Products*, 75 (2), pp 175–180.
- Tanada, Y. and H. K., Kayea. (1993). *Insect pathology, "fungal infection"*. Academic Press, San Diego. Pp 321-387.
- Tobe S. S. and Pratt G. E. (1975). Corpus allatum activity in vitro during ovarian maturation in the desert locust, *Schistocerca gregaria*. *Journal of Experimental Biology*, 62, 611-627.
- Wafford, K. A. and D. B., Sattelle (1989). L-glutamate receptors on the cell body membrane of an identified insect motor neurone. *J. Exp. Biol.* 144: 449-462.
- Wolstenholme A.J. and A. T. Rogers (2005). Glutamate-gated chloride channels and the mode of action of the avermectin/milbemycin anthelmintics. *Parasitology* 131:585–595.
- Woo, W.S.; Chi, H.J.; Yun and Hye, S. (1977). Alkaloid screening of some Saudi Arabian plants. *Saengyak Hakhoe Chi (Hanguk Saengyak Hakhoe)*, 8(3): 109-113.

## الاستجابات الفسيولوجية وتحت تركيبه للجراد الصحراوي عند معاملة بفطره الميتاريزيم اكريديوم واربع مركبات ذات نشاط حيوي أيمن فراج أحمد<sup>١</sup> و ثروت عبد المنعم عبد الفتاح<sup>٢</sup> و جمال محمد عبد اللطيف<sup>٢</sup> و محمد خيرى الديقاموني<sup>٢</sup>

١ جامعة الأزهر، كلية العلوم، قسم الميكروبيولوجي، مدينة نصر، القاهرة  
٢ معهد بحوث وقاية النباتات، مركز البحوث الزراعي، الدقى، جيزه

تم دراسة تأثير فطر الميتاريزيم اكريديوم وكذلك ٤ مركبات نشطه حيويًا (النييم و الحمض الأمين جلوتاميك ومستخلص نبات الشينيس مولى و الأباكتين) بعشر الجرعه المقرره منفرده او مخلوطه مع الفطر على نسبة الموت والتغيرات الهستولوجيه الدقيقه لأنسجه المبيض لحوريات العمر الخامس للجراد الصحراوي واطهرت النتائج ان نسبة الموت فى حاله فطر الميتاريزيم انيسوبلى وصلت الى ١٠٠% بعد ١٤ يوم اما عند استخدام النيم فكانت نسبة الموت ٦٠% بعد ١٢ يوم وعند استخدام الحمض الأمينى جلوتاميك فكانت نسبة الموت ٢٠% بعد ١٠ ايام وعند استخدام مستخلص من نبات الشينيس مولى اصبحت نسبة الموت ٥٠% بعد ١٤ يوم اما عند استخدام الأباكتين فكانت نسبة الموت ٦٠% بعد ١١ يوم بينما جميع المخاليط ادت الى ١٠٠% موت بعد ٥ ايام. بينما عند دراسته تأثير المواد سالفه الذكر فى الحاله المنفرده على المده الزمنيه اللازمه لاحداث نسب موت ٢٥%, ٥٠% و ٩٠% لحشره الجراد الصحراوي وجد انها تستغرق وقت طويل وكان فطر الميتاريزيم اشدها تأثيرا حيث وجد ان المده الزمنيه اللازمه لاحداث نسبة ٥٠% موت هي ٧،٢٦١ يوم، بينما عند خلط فطر الميتاريزيم مع المواد الاخرى كلا على حده فانها تقوم بتسريع عمليه الموت بصوره ملحوظه وكان اسرع المخاليط موتا مخلوط فطر الميتاريزيم مع النيم حيث كانت المده الزمنيه اللازمه لاحداث ٥٠% موت هي ١،٦٢٥ يوم.

وعند دراسته تأثير المواد السابقه على التغيرات الواقعه على مبيض حشره الجراد ودراسه التأثير بواسطه استخدام الميكروسكوب الالكترونى النافذ فوجد ان هناك تغيرات حدثت وان تلك التغيرات تنوعت واختلفت من معاملة الى اخرى وشملت انفجار النواه، تغير فى شكل النواه، تجمع الكروماتين، وجود فجوات، تغير فى شكل وحجم الميتوكوندريا، ظهور قطرات زيتيه وفى المعاملات التى تحتوى على فطر الميتاريزيم انيسوبلى فان الخيوط الفطريه كانت ظاهره جليا فى مبيض الحشره.