

## AN EVALUATION OF TWO TOMATO CULTIVARS TO INFESTATION BY CERTAIN INSECT PESTS

Abdallah, Y. E. Y. and Hana S. A. Faraj  
Plant Protection Dept., Fac. Agric., Ain Shams Univ., Cairo,  
Egypt.



### ABSTRACT

Experiments were carried out at tomato field at Qalyubia Governorate, Egypt cultivated with two different cultivars to test its susceptibility to infestation with *Bemisia tabaci* (Gennadius); aphids {*Aphis gossypii* and *Myzus persicae* (Sülzer)}; *Empoasca decipiens* Paoli and *Tuta absoluta* (Meyrick) in addition to the seasonal abundance of these pests on the two cultivars in two successive summer seasons (2012 and 2013). Results indicated that *B. tabaci* and aphids had two peaks of abundance in May and June in both years on both cultivars. *E. decipiens* also had two peaks of abundance in May and June in both years with no significant difference between the two cultivars. The two tested cultivars almost harbored the same population density of the immature stages of *T. absoluta* during summer season, 2012. The pest had only one peak during the last week of May. There was negative and highly significant correlations between the population densities of all considered pests and the numbers of hairs (trichomes) present on tomato leaves sampled on 20 May and 4 June except for aphids which disappeared during hot summer months on the cultivar Hybrid Super. In the third sampling date (20 June, 2013), the correlation was also negative and highly significant with the population of *B. tabaci* and positive with the population of *T. absoluta*. The same trend was observed on the cultivar Crystal HYB. The effects of (N), (P) and (Ca) were positive and highly significant on all considered pests on both cultivars. The effect of (K) was negative and highly significant on all considered pests on both cultivars. The effect of (Mg) was positive and highly significant on the cultivar Hybrid; while it was negatively correlated with sap sucking pests only on the cultivar Crystal. Data indicate that the effect of Fe, Zn and Mn was positive and highly significant on both cultivars and Cu was negatively affected these pests on both cultivars.

### INTRODUCTION

Tomato (*Lycopersicon esculentum* L.) is one of the most popular vegetable crops in the world (Mukhtar *et al.*, 2009) mainly grown for human consumption. In 2012, the total area grown with tomatoes in the world was about 11.2 million feddans and the total world production was 161,793,834 tons. Egypt is ranked as the fifth country in the world in terms of total production of tomatoes. The cultivated area reached 591,384 feddans produced 8,625,219 tons (Food and Agriculture Organization Statistics, 2014).

Tomato plants and fruits are attacked by many insect pests in the field. The most important insect pests are tomato whitefly, *Bemisia tabaci* (Gennadius); aphids {cotton aphid, *Aphis gossypii* (Glover) and green peach aphid, *Myzus persicae* (Sülzer)}; American tomato budworm, *Helicoverpa armigera* (Hübner) and tomato leafminer, *Tuta absoluta* (Meyrick) which are considered the most destructive pests of tomato yield all over the world in addition to potato leafhopper, *Empoasca decipiens* Paoli. The ecology of

these pests was studied by Arnal *et al.*, 1993; Adam *et al.*, 1997; Bezerra *et al.*, 2004; Garzia *et al.*, 2009; Setiawati *et al.*, 2009; Summers *et al.*, 2010; Chakraborty, 2011; Hrcic and Radoniic 2011; Allache and Demnati, 2012 and Spasov *et al.*, 2013. The susceptibility of different cultivars was also studied by (Leite *et al.*, 1999; Abdallah *et al.*, 2001; Leite *et al.*, 2004; Marcos *et al.*, 2005; Sahu and Shaw, 2005; Ashfaq *et al.*, 2012; Snyder and Min, 2012 and Han *et al.*, 2014). The present investigation was conducted to follow up the seasonal fluctuations in the population densities of *B. tabaci*, aphids, *E. decipiens* and *T. absoluta* on tomato plants and to investigate the susceptibility of two grown cultivars to the infestation with these pests.

## **MATERIALS AND METHODS**

Field experiments were carried out at the experimental farm attached to the Faculty of Agriculture, Ain Shams University at Shalakan, Qalyubya Governorate. An area of about 0.25 feddan was cultivated with each one of the two tested tomato cultivars in spring-summer plantations 2012 and 2013 (Hybrid Super Strain BF<sub>1</sub> and Super Crystal HYB). Transplanting dates were 4 April, 2012 and 24 March, 2013. The agricultural practices were carried out as usual. The seasonal abundance and population dynamics of tomato pests viz.; tomato whitefly, *Bemisia tabaci* (Gennadius), aphids (cotton aphid, *Aphis gossypii* (Glover) and green peach aphid, *Myzus persicae* (Sülzer)), potato leafhopper, *Empoasca decipiens* Paoli and tomato leafminer, *Tuta absoluta* were studied.

### **Population dynamics of tomato pests**

The seasonal abundance of different stages of *B. tabaci*, (adults, nymphs and eggs) were counted on 20 leaflets, replicated three times (a total of 60 leaflets/ sampling date). During the first plantation, sampling period extended from 8 May to 10 July, 2012 (10 inspections). In the second plantation, this period extended from 23 April to 2 July, 2013 (11 inspections). This procedure was carried out for each tested tomato cultivar. Counts of aphids and leafhoppers (adults and nymphs) were carried out at the same way. For *T. absoluta*, numbers of eggs, larvae and mines were counted. The total number for each insect pest was used to represent its population dynamics.

Susceptibility of two tomato cultivars to pest infestation

### **Anatomical studies**

Anatomical studies were carried out where leaf samples were picked up from the upper part of the plant from the two tested cultivars at three sampling dates, 12 May; 4 June and 20 June, 2013. Samples were killed and fixed in FAA using the method of Jonhnsen (1940), dehydrated with ascending concentrations of ethyl alcohol and serial transverse sections (20µ each) were prepared using the standard paraffin technique. All sections were fixed in small glass slides (1x2 cm) by means of Haupt's adhesive. Finally, specimens were sputter coated with gold coat after removing wax. To study the number and length of trichomes (hairs) on both sides of the leaves, specimens (0.2x0.8 mm) were killed and fixed in glutaraldehyde 2.5% for 24h

at 4°C, then post-fixed in osmium tetroxide (O<sub>8</sub>O<sub>4</sub>) 1% for one hour at room temperature (Millonig, 1961), dehydrated with ascending concentrations of acetone. Samples were critical point dried and coated with gold. All observations, measurements and photography were done through a Joel Scanning Electron Microscope (SEM) (T. 330A) at 30 Kv. at the Unit of Electron Microscopy, Central Laboratory Unit, Faculty of Agriculture, Ain Shams University.

#### **Macro- and micro-nutrients**

Leaf samples from the two tested tomato cultivars were taken for nutritional studies at the same dates mentioned above. Leaves were dried at 70°C, weighted and digested using a mixture of sulfuric acid with hydrogen peroxide according to the method described by FAO (1989). Wet digestion was performed through using 0.5 g from oven-dry leaves material and then added in 50 ml conical flask and digested with 10 ml of H<sub>2</sub>SO<sub>4</sub> on a hot plate at approximately 270°C. Small quantities of H<sub>2</sub>O<sub>2</sub> were added repeatedly until the digested solution became clear. The solution was left to cool and then diluted to 50 ml with redistilled water in a volumetric flask.

In the acid digested solution, the micronutrients determined were iron (Fe), Zinc (Zn), manganese (Mn) and copper (Cu) using atomic absorption spectrophotometer, while Phosphorus (P) concentration in acid digested samples was determined by colorimetric method (ammonium molybdate) using spectrophotometer. Total nitrogen content (N) was determined using microkjeldahle method, whereas Potassium (K) content was detected using the flame photometer (Chapman and Pratt, 1982). All nutritional studies were conducted at the Arid Land Agricultural Research and Services Center, Faculty of Agriculture, Ain Shams University.

## **RESULTS AND DISCUSSION**

### **The fluctuations in the population densities of certain tomato pests on two tomato cultivars during summer plantations**

#### **A) *Bemisia tabaci* (Gennadius)**

Data in Table (1) show the fluctuations in the population densities of the four investigated tomato pests on two cultivars of tomato during summer plantation of 2012. These data indicate that on the cultivar Hybrid Super, the population of *B. tabaci* had two peaks of abundance during that season; the first was during the second week of May (1227 individuals/60 leaflets) compared to 405 individuals in the second peak at the second week of June. A gradual decrease in the population density occurred till the end of the season. On the cultivar Crystal HYB, the total population of *B. tabaci* was less than that on Hybrid Super and also had two peaks of abundance; the first was during the second half of May (807 individuals) and the second was one month later (388 individuals). The differences between the two cultivars in their infestation were insignificant ( $t = 2.02$ ).

Data in Table (2) clearly indicate that the population of *B. tabaci* during 2013 plantation was higher than that of summer 2012. On the cultivar Hybrid Super, the total population reached 6286 individuals with three peaks of

abundance. The first peak occurred during the last week of April, 2013 being 1803 individuals. The second occurred during the second week of May, 2013 (946 individuals); while the third peak was during the first week of June (318 individuals). On the cultivar Crystal HYB, the population of *B. tabaci* was much lower (3673 individuals) than that on Hybrid and also had three peaks of abundance; the first was during the last week of April, 2013 with a total of 1045 individuals/60 leaflets. The second peak was reached two weeks later with a total of 603 individuals; while the third was during the first week of June being 226 individuals. The differences between the two cultivars in their infestation were significant ( $t = 2.20^*$ ).

**B) Aphids, *Aphis gossypii* (Glover) and *Myzus persicae* (Sülzer)**

Regarding the population size of aphids occurred on the two tested cultivars, the case was on the contrary of that of *B. tabaci* since the cultivar Hybrid Super harbored a relatively lower population than Crystal HYB during 2012 summer season ( $t = 0.61$ ). Two peaks of abundance were observed on each cultivar during the second week of May and one month later. During 2013 season, the same trend was observed and although the differences between the population densities on the two cultivars were higher but it was insignificant ( $t = 1.02$ ). Only one peak was observed on each cultivar at the beginning of the season (last week of April).

**Table 1: The total numbers of certain tomato pests on two cultivars of tomato during summer plantation (2012), Shalakan, Qalyubya Governorate**

Sampling Date	Planting date 4-4-2012							
	Hybrid Super cultivar				Crystal H Y B cultivar			
	<i>B. tabaci</i>	Aphids	<i>E. decipiens</i>	<i>T. absoluta</i>	<i>B. tabaci</i>	Aphids	<i>E. decipiens</i>	<i>T. absoluta</i>
08/5	1227	330	21	19	794	477	19	25
15/5	930	122	95	57	807	110	89	31
22/5	805	42	68	57	692	9	31	19
29/5	507	7	57	80	530	3	70	72
05/6	391	9	12	27	263	5	9	27
12/6	405	12	14	33	388	15	17	23
19/6	213	1	23	15	167	0	23	9
26/6	115	0	2	3	86	1	0	6
03/7	33	0	0	2	41	0	0	4
10/7	3	0	0	1	3	0	0	1
Total	4629	523	292	294	3771	620	258	217
't' value	2.02	0.61	0.84	1.73				

**C) Potato leafhopper, *Empoasca decipiens* Poali**

The population of *E. decipiens* in 2012 season was higher than that of 2013. During 2012 season, the population had two peaks of abundance on both cultivars; the first was during the second week of May and the second was during the third week of June. These peaks reached 95 and 23 leafhoppers/60 leaflets on Hybrid Super and 89 and 23 leafhoppers/60 leaflets on Crystal HYB. The differences between the total populations on

both cultivars were insignificant;  $t = 0.84$  (Table 1). In the second season (2013), the population of *E. decipiens* was much lower than aphids on both cultivars and it had only one peak on the second week of May (74 leafhoppers/60 leaflets) on Hybrid Super and 63 leafhoppers on Crystal HYB. The differences between the populations on both cultivars were insignificant;  $t = 0.64$  (Table 2).

**D) Tomato leafminer, *Tuta absoluta* (Meyrick)**

Data in Tables (1 & 2) reveal that the two tested cultivars almost harbored the same population density of the immature stages of *T. absoluta* during summer season 2012. The differences in the population density were insignificant ( $t = 1.73$ ). The pest had only one peak during the last week of May in 2012 being 80 and 72 individuals/60 leaflets on Hybrid Super and Crystal HYB, respectively. This peak occurred on the first week of May in 2013 being 66 and 46 individuals on both cultivars, respectively. The differences in the population density were insignificant ( $t = 2.26^*$ ).

**Table 2: The total numbers of certain tomato pests on two cultivars of tomato during summer plantation (2013), Shalakan, Qalyubya Governorate**

Sampling Date	Planting date 24-3-2013							
	Hybrid Super cultivar				Crystal H Y B cultivar			
	<i>B. tabaci</i>	Aphids	<i>E. decipiens</i>	<i>T. absoluta</i>	<i>B. tabaci</i>	Aphids	<i>E. decipiens</i>	<i>T. absoluta</i>
23/4	1295	641	18	4	691	1006	8	6
30/4	1803	180	21	31	1045	227	30	31
07/5	903	53	47	66	495	51	63	46
14/5	946	35	74	47	603	12	28	26
21/5	399	0	3	23	191	0	3	13
28/5	212	0	4	12	217	0	4	10
04/6	318	0	2	8	226	0	0	0
11/6	211	0	0	18	110	0	0	9
18/6	110	0	0	5	94	0	0	7
25/6	76	0	0	0	45	0	0	0
02/7	13	0	0	0	19	0	0	2
Total	6286	909	169	214	3736	1296	136	150
t' value	2.2*	1.06	0.64	2.26*				

Most authors in the literature had results that not in full harmony with our results. Arno *et al.* (2005), WenFeng *et al.* (2009) and Karut *et al.* (2012) reported that *B. tabaci* was more abundant in autumn than in spring and its populations were strongly reduced during winter. These results agreed with those found by Herakly (1974) in Egypt who found that *Myzus persicae* and *Aphis gossypii* infested tomato plants was found to be higher in late summer season. Dawood (1999) recorded one population peak of *A. gossypii* on the 18 and 21 May during two seasons on the five tomato cultivars. Abdallah *et al.* (2000) reported that *M. persicae* was detected during summer plantation of 1995 on potato plants in reliable numbers on the 4<sup>th</sup> week of April when plants were 7 weeks old. In 1996, this peak took place one week earlier. Similar results were found by Vuong *et al.* (2001); Ebadah (2002) and Der *et*

al. (2003). These data are also in harmony with those found by Oliveira *et al.* (2008) in Brazil; Cota (2011); Nannini *et al.* (2011), Hrcic and Radonić (2011) and Allache and Demnati (2012) in Algeria.

**The relationship between the numbers of hairs on tomato leaves and population densities of the considered pests**

Data in Table (3) show the correlation between the number of hairs (at different magnification powers of SEM) in the cultivar Hybrid sown during summer 2013 and the population densities of certain tomato pests. These data reveal the presence of negative and highly significant correlation between the population densities of all considered pests and the numbers of hairs during the first sampling date (20 May). The correlation coefficient values were the same during the second sampling date (4 June) except for aphids which disappeared during hot summer months. In the third sampling date (20 June), the correlation was also negative and highly significant with the population of *B. tabaci* and positive with the population of *T. absoluta*. On the cultivar Crystal (Table 4), the correlation was also negative and highly significant between *B. tabaci* populations and the numbers of hairs during the three sampling dates; while the correlation with *T. absoluta* was positive and insignificant.

**Table 3: The correlation coefficient values for the relationship between the population densities of certain pests and number of hairs on tomato leaflets (cultivar Hybrid) during spring-summer plantation of 2013**

Pest	Population density	Average no. of hairs/10 mm <sup>2</sup> at		
		150X	200X	500X
First Sampling Date (12/5/2013)				
No. of hairs		(500)	(500)	(150)
<i>Bemisia tabaci</i>	946	-0.9780	-0.7502	-0.8457
Aphids	35	-0.7890	-0.9875	-0.7882
<i>E. decipiens</i>	74	-0.8894	-0.7928	-0.9171
<i>Tuta absoluta</i>	47	-0.9632	-0.9338	-0.8817
Second Sampling Date (4/6/2013)				
No. of hairs		(600)	(450)	(150)
<i>Bemisia tabaci</i>	318	-0.9632	-0.9781	-0.9359
Aphids	0	0.000	0.000	0.000
<i>E. decipiens</i>	2	-0.9231	-0.7645	-0.8825
<i>Tuta absoluta</i>	8	0.8704	0.8191	-0.9266
Third Sampling Date (20/6/2013)				
No. of hairs		(366)	(577)	(416)
<i>Bemisia tabaci</i>	110	-0.9618	-0.8952	-0.8972
Aphids	0	0.000	0.000	0.000
<i>E. decipiens</i>	0	0.000	0.000	0.000
<i>Tuta absoluta</i>	5	0.2092	0.5378	0.1353

The obtained results are in agreement with those found by Zareh (1987) who found insignificant correlation between infestation by *Empoasca* spp. and the density of hairs and glands on lower surface of cotton leaf. Amr (1993) found a negative correlation between jassid infestation on different cotton varieties and the number of stellate and glandular trichomes in leaf

epidermis. Heinz and Zalom (1995) who mentioned that tomato cultivars with low trichome densities sustained less whitefly oviposition. Muigai *et al.* (2002) found that most dead adults of *B. argentifolii* were trapped in glandular trichome exudates. Mohasin *et al.* (2005) reported that the increase in the number of leaf hair per unit area reduced the population density of *B. tabaci*. Sahu and Shaw (2005) reported that the presence of trichomes on tomato plants had negative effect on whitefly incidence. Also, Samarajeewa *et al.* (2005) found that tomato plants resistant to infestation with *B. tabaci* had more trichomes on leaves and stems than the susceptible plants. JiRong *et al.* (2011) reported that the hairy tomato plants can repel aphids, *B. tabaci* and American leaf miner. Snyder and Min (2012) found that tomato accessions resistant to aphids were also resistant to *Liriomyza* sp.

**Table 4: The correlation coefficient values for the relationship between the population densities of certain pests and number of hairs on tomato leaflets (cultivar Crystal) during spring-summer plantation of 2013.**

Pest	Population density	Average no. of hairs/10 mm <sup>2</sup> at		
		150X	200X	500X
First Sampling Date (12/5/2013)				
No. of hairs		(250)	(250)	(150)
<i>Bemisia tabaci</i>	603	-0.9660	-0.9478	-0.9526
Aphids	12	-0.9181	-0.9111	-0.9358
<i>E. decipiens</i>	28	-0.8161	-0.8086	-0.7993
<i>Tuta absoluta</i>	26	0.2055	0.1347	0.1229
Second Sampling Date (4/6/2013)				
No. of hairs		(500)	(450)	(150)
<i>Bemisia tabaci</i>	226	-0.8980	-0.9662	-0.9091
Aphids	0	0.000	0.000	0.000
<i>E. decipiens</i>	0	0.000	0.000	0.000
<i>Tuta absoluta</i>	0	0.000	0.000	0.000
Third Sampling Date (20/6/2013)				
No. of hairs		(569)	(399)	(102)
<i>Bemisia tabaci</i>	94	-0.9911	-0.9924	-0.9590
Aphids	0	0.000	0.000	0.000
<i>E. decipiens</i>	0	0.000	0.000	0.000
<i>Tuta absoluta</i>	7	0.2713	0.2669	0.0952

**The effect of major nutritional components in tomato leaves on the population densities of the considered pests**

Data in Tables (5 & 6) reveal that the effect of (N) was positive and highly significant on all considered pests on the cultivars Hybrid Super and Crystal HYB. The effect of (P) was positive and highly significant on all sap sucking pests on the cultivar Crystal, the same relation was found with *T. absoluta* on the cultivar Hybrid. The effect of (K) was negative and highly significant on all considered pests on both cultivars. The effect of (Ca) was positive and highly significant on all considered pests on both cultivars. The effect of (Mg) was positive and highly significant on the cultivar Hybrid; while it was significantly correlated with sap sucking pests only on the cultivar Crystal.

**Table 5: The correlation coefficient values for the relationship between the population densities of certain pests and major nutritional components in tomato leaflets (cultivar Hybrid) during spring-summer plantation of 2013**

Sample	<i>B. tabaci</i>	N%	"r"	Aphids	N %	"r"	<i>E. decipiens</i>	N %	"r"	Tuta	N %	"r"
1	1803	3.87	0.8148	180	3.87	0.7119	21	3.87	0.7984	31	3.87	0.9993
2	399	3.70		0	3.70		3	3.70		23	3.70	
3	110	3.25		0	3.25		0	3.25		5	3.25	
	<i>B. tabaci</i>	P%	"r"	Aphids	P %	"r"	<i>E. decipiens</i>	P %	"r"	Tuta	P %	"r"
1	1803	0.55	0.0524	180	0.55	-0.1076	21	0.55	0.0246	31	0.55	0.5926
2	399	0.93		0	0.93		3	0.93		23	0.93	
3	110	0.29		0	0.29		0	0.29		5	0.29	
	<i>B. tabaci</i>	K%	"r"	Aphids	K %	"r"	<i>E. decipiens</i>	K %	"r"	Tuta	K %	"r"
1	1803	0.86	-0.7060	180	0.86	-0.5839	21	0.86	-0.6861	31	0.86	-0.9791
2	399	1.17		0	1.17		3	1.17		23	1.17	
3	110	3.69		0	3.69		0	3.69		5	3.69	
	<i>B. tabaci</i>	Ca%	"r"	Aphids	Ca %	"r"	<i>E. decipiens</i>	Ca %	"r"	Tuta	Ca %	"r"
1	1803	5.22	0.9521	180	5.22	0.8910	21	5.22	0.9432	31	5.22	0.9635
2	399	3.36		0	3.36		3	3.36		23	3.36	
3	110	1.81		0	1.81		0	1.81		5	1.81	
	<i>B. tabaci</i>	Mg%	"r"	Aphids	Mg %	"r"	<i>E. decipiens</i>	Mg %	"r"	Tuta	Mg %	"r"
1	1803	0.80	0.8846	180	0.80	0.7988	21	0.80	0.8713	31	0.80	0.9953
2	399	0.67		0	0.67		3	0.67		23	0.67	
3	110	0.47		0	0.47		0	0.47		5	0.47	

**Table 6: The correlation coefficient values for the relationship between the population densities of certain pests and major nutritional components in tomato leaflets (cultivar Chrystal) during spring-summer plantation of 2013**

Sample	<i>B. tabaci</i>	N%	"r"	Aphids	N %	"r"	<i>E. decipiens</i>	N %	"r"	Tuta	N %	"r"
1	1045	4.25	0.5380	227	4.25	0.4576	30	4.25	0.5364	31	4.25	0.8597
2	191	4.30		0	4.30		3	4.30		13	4.30	
3	94	3.38		0	3.38		0	3.38		7	3.38	
	<i>B. tabaci</i>	P%	"r"	Aphids	P %	"r"	<i>E. decipiens</i>	P %	"r"	Tuta	P %	"r"
1	1045	0.61	0.7792	227	0.61	0.7178	30	0.61	0.7781	31	0.61	0.1343
2	191	0.50		0	0.50		3	0.50		13	0.50	
3	94	0.22		0	0.22		0	0.22		7	0.22	
	<i>B. tabaci</i>	K%	"r"	Aphids	K %	"r"	<i>E. decipiens</i>	K %	"r"	Tuta	K %	"r"
1	1045	1.24	-0.6292	227	1.24	-0.5544	30	1.24	-0.6277	31	1.24	-0.7619
2	191	1.43		0	1.43		3	1.43		13	1.43	
3	94	3.90		0	3.90		0	3.90		7	3.90	
	<i>B. tabaci</i>	Ca%	"r"	Aphids	Ca %	"r"	<i>E. decipiens</i>	Ca %	"r"	Tuta	Ca %	"r"
1	1045	4.61	0.8017	227	4.61	0.7429	30	4.61	0.8006	31	4.61	0.9739
2	191	3.84		0	3.84		3	3.84		13	3.84	
3	94	2.17		0	2.17		0	2.17		7	2.17	
	<i>B. tabaci</i>	Mg%	"r"	Aphids	Mg %	"r"	<i>E. decipiens</i>	Mg %	"r"	Tuta	Mg %	"r"
1	1045	0.62	-0.2863	227	0.62	-0.3738	30	0.62	-0.2881	31	0.62	0.9199
2	191	1.07		0	1.07		3	1.07		13	1.07	
3	94	0.54		0	0.54		0	0.54		7	0.54	

**The relationship between the minor nutritional components in tomato leaves and population densities of the considered pests**

Data in Tables (7 & 8) show the correlation between the percentage of minor nutritional components in tomato leaves sown during summer 2013 and the population densities of the considered tomato pests. These data clearly indicate that the effect of the minor nutrients Fe, Zn and Mn was positive and highly significant on both cultivars and Cu was negatively affected these pests on both cultivars. Again the results of the summer plantation proved to be more reliable in expressing the effect of both major and minor nutrients.

**Table 7: The correlation coefficient values for the relationship between the population densities of certain pests and minor nutritional components in tomato leaflets (cultivar Hybrid) during spring-summer plantation of 2013**

Sample	<i>B. tabaci</i>	Fe ppm	"r"	Aphids	Fe ppm	"r"	<i>E. decipiens</i>	Fe ppm	"r"	<i>Tuta</i>	Fe ppm	"r"
1	1803	6400		180	6400		21	6400		31	6400	
2	399	2300	0.9491	0	2300	0.9872	3	2300	0.9575	23	2300	0.6197
3	110	3000		0	3000		0	3000		5	3000	
	<i>B. tabaci</i>	Zn ppm	"r"	Aphids	Zn ppm	"r"	<i>E. decipiens</i>	Zn ppm	"r"	<i>Tuta</i>	Zn ppm	"r"
1	1803	41.75		180	41.75		21	41.75		31	41.75	
2	399	34.50	0.8752	0	34.50	0.7868	3	34.50	0.8615	23	34.50	0.9970
3	110	22.50		0	22.50		0	22.50		5	22.50	
	<i>B. tabaci</i>	Mn ppm	"r"	Aphids	Mn ppm	"r"	<i>E. decipiens</i>	Mn ppm	"r"	<i>Tuta</i>	Mn ppm	"r"
1	1803	127.25		180	127.25		21	127.25		31	127.25	
2	399	93.00	0.9028	0	93.00	0.8226	3	93.00	0.8905	23	93.00	0.9906
3	110	47.50		0	47.50		0	47.50		5	47.50	
	<i>B. tabaci</i>	Cu ppm	"r"	Aphids	Cu ppm	"r"	<i>E. decipiens</i>	Cu ppm	"r"	<i>Tuta</i>	Cu ppm	"r"
1	1803	47.50		180	47.50		21	47.50		31	47.50	
2	399	84.25	0.0085	0	84.25	-0.1680	3	84.25	-0.0363	23	84.25	0.5424
3	110	27.50		0	27.50		0	27.50		5	27.50	

Natarajan (1987) observed that the population of *B. tabaci* escalated as the level of nitrogen increased, but decreased with increased level of potassium with the same level of phosphorus. Purohit and Deshpande (1991) proved that normal and double rates of NPK fertilizer increased *B. tabaci* infestation of cotton plants compared with untreated plants. Rote and Puri (1992) found a highly significant positive correlation between the population of *B. tabaci* and on cotton plants nitrogen content of leaves. Abdallah *et al.* (2001) found that N and K contents in cotton leaves were negatively correlated with populations of *A. gossypii*, *Empoasca* sp. and *B. tabaci*. They added that all pests' populations were positively correlated with Zn, Fe and Mg content. Feltrin *et al.* (2004) reported that potassium source had no significant effect on the number of eggs and nymphs of *B. tabaci*. Leite *et al.* (2004) reported that N and K contents in tomato leaves had no effect on the

egg population of *T. absoluta*. Again Leite *et al.* (2004) found that N and K contents in tomato leaves had insignificant effect on the population of *B. tabaci*. Ashfaq *et al.* (2012) reported that ferrous (Fe<sup>2+</sup>) and phosphorous content in tomato leaves were negatively correlated with larval population of *Helicoverpa armigera*; whereas nitrogen, calcium, magnesium, manganese and zinc content were positively correlated with larval population. Han *et al.* (2014) reported that the development time for *T. absoluta* from egg to adult was negatively correlated with tomato leaf N.

**Table 8: The correlation coefficient values for the relationship between the population densities of certain pests and minor nutritional components in tomato leaflets (cultivar Crystal) during spring-summer plantation of 2013**

Sample	<i>B. tabaci</i>	Fe ppm	"r"	Aphids	Fe ppm	"r"	<i>E. decipiens</i>	Fe ppm	"r"	<i>Tuta</i>	Fe ppm	"r"
1	1045	6300		227	6300		30	6300		31	6300	
2	191	5600	0.7943	0	5600	0.7346	3	5600	0.7931	13	5600	0.8760
3	94	4000		0	4000		0	4000		7	4000	
	<i>B. tabaci</i>	Zn ppm	"r"	Aphids	Zn ppm	"r"	<i>E. decipiens</i>	Zn ppm	"r"	<i>Tuta</i>	Zn ppm	"r"
1	1045	41.25		227	41.25		30	41.25		31	41.25	
2	191	40.75	0.5930	0	40.75	0.5159	3	40.75	0.5915	13	40.75	0.7065
3	94	17.5		0	17.50		0	17.50		7	17.50	
	<i>B. tabaci</i>	Mn ppm	"r"	Aphids	Mn ppm	"r"	<i>E. decipiens</i>	Mn ppm	"r"	<i>Tuta</i>	Mn ppm	"r"
1	1045	115.25		227	115.25		30	115.25		31	115.25	
2	191	94.50	0.7956	0	94.50	0.7360	3	94.50	0.7944	13	94.50	0.8771
3	94	47.50		0	47.50		0	47.50		7	47.50	
	<i>B. tabaci</i>	Cu ppm	"r"	Aphids	Cu ppm	"r"	<i>E. decipiens</i>	Cu ppm	"r"	<i>Tuta</i>	Cu ppm	"r"
1	1045	32.50		227	32.5		30	32.5		31	32.50	
2	191	151.73	0.3850	0	151.73	0.4689	3	151.73	0.3868	13	151.73	0.2430
3	94	27.50		0	27.50		0	27.5		7	27.50	

## REFERENCES

- Abdallah, Y. E. Y.; M. A. Daoud and G. B. El-Saadany (2000). Seasonal abundance and spatial distribution pattern of *Myzus persicae* Sulzer in potato field. *Adv. Agric. Res.* 5 (3): 1557-1574.
- Abdallah, Y. E. Y.; S. I. Ibrahim; E. M. Abd-Elmoniem and L. A. Youssef (2001). Incidence of some piercing-sucking insects in relation to morphological leaf characters, some chemical and nutritional components of some cotton cultivars. *Annals Agric. Sci., Ain Shams Univ., Cairo*, 46 (2): 807-827.
- Adam, M. K.; M. A. Bachatly and S. A. Doss (1997). Population of the whitefly *Bemisia tabaci* (Genn.) (Homoptera: Aleyrodidae) and its parasitoid *Eretmocerus mundus* Mercet (Hymenoptera : Aphelinidae) in protected cucumber cultivations. *Egypt. J. Agric. Res.*, 75 (4): 939-950.

- Allache, F. and Demnati, Fatma (2012). Population changes of *Tuta absoluta* (Mey.) (Lepidoptera, Gelichiidae): A new introduced tomato crop pest at Biskra in Algeria. *Jordan J. Agric. Sci.*, 8 (3): 391-400.
- Arnal, E.; E. Debrot; R. Marcano and A. Montagne (1993). Population fluctuation of whiteflies and its relation to tomato yellow mosaic in one location in Venezuela. *Fitopatologia Venezolana*, 6 (1): 21-26.
- Arno, J.; M. Matas; M. Marti; J. Arino; J. Roig and R. Gabarra (2005). Coexistence between *Trialeurodes vaporariorum* and *Bemisia tabaci* and impact of natural enemies in tomato crops under Mediterranean conditions. *Bulletin OILB/ SROP*, 28 (1): 1-4.
- Ashfaq, M.; M. Sajjad; M. N. Ane and N. Rana (2012). Morphological and chemical characteristics of tomato foliage as mechanisms of resistance to *Helicoverpa armigera* (Hübner) (Lepidoptera:Noctuidae) larvae. *African J. Biotech.*, 11 (30): 7744-7750.
- Bezerra, Mary-Ann, S.; Maria, R. V. de Oliveira and S. D. Vasconcelos (2004). Does the presence of weeds affect *Bemisia tabaci* (Gennadius) (Hemiptera:Aleyrodidae) infestation on tomato plants in a semi-arid groecosystem?. *Neotrop. Entomol.*, 33 (6): 769-775.
- Chakraborty, K. (2011). Incidence of Aphid, *Aphis gossypii* Glover (Hemiptera: Aphidae) on tomato crop in the agro climatic conditions of the northern parts of West Bengal, India. *World J. Zool.*, 6 (2): 187-191.
- Chapman, H. D. and P. F. Pratt (1982). *Methods of analysis for soil, plant and waters*. Priced publication 4034, Univ. Calif. Division of Agric. Sci.
- Cota, E. (2011). IPM of aphids in vegetable field crops in Albania. *IOBC/WPRS Bulletin*. 65: 13-16.
- Dawood, M. Z. (1999). Susceptibility of different tomato cultivars to infestation with *Aphis gossypii* Glover at Damietta Governorate. *Egypt. J. Agric. Res.*; 77 (4): 1645-1655.
- Der, Z.; B. Penzes and A. Orosz (2003). Species Composition of leafhoppers collected in tomato fields. *Noveny Vedelem*. 39 (10): 485-494.
- Ebadah, I. M. A. (2002). Population fluctuations and diurnal activity of the leafhopper, *Empoasca decipiens* on some summer crops in Kalubia Governorate, Egypt. *Bull. Fac. Agric., Cairo Univ.*; 53 (4): 653-670.
- FAO Soil Bulletin (1989). *Soil and plant testing*. 38 (2): 250pp.
- FAO (2014). *Food and Agriculture Organization Statistics*, 110 pp.
- Garzia, G. T.; G. Siscaro; A. Colombo and G. Campo (2009). Reappearance of *Tuta absoluta* in Sicily. *Italian Informatore Agrario*, 65 (4): 71.
- Han, P.; A. V. Lavoit; J. L. Bot; E. A. Desneux and N. Desneux (2014). Nitrogen and water availability to tomato plants triggers bottom-up effects on the leafminer, *Tuta absoluta*. *Scientific Reports* 4; Article number: 4455, doi:10.1038/ srep04455.
- Heinz, K. M. and F. G. Zalom (1995). Variation in trichome-based resistance to *Bemisia argentifolii* (Homoptera: Aleyrodidae) oviposition on tomato. *J. Econ. Entom.*, 88 (5): 1494-1502.
- Herakly, F. A. (1974). Preliminary survey of pests infesting solanaceous truck crops in Egypt. *Bull. Ent. Soc. Egypte*. 47: 2133-2140.

- Hrnčić, S. and S. Radonić (2011). *Tuta absoluta* Meyrick (Lepidoptera: Gelechiidae) a new pest in Montenegro. IOBC/WPRS Bulletin, 68:71-74.
- Ji Rong, Z.; W. Hui Li and W. Jia Ming (2011). Breeding of 'Hangza No.401'- a new tomato cultivar with good quality and disease resistance. J. Anhui Agric. Univ., 38 (1): 110-117.
- Johansen, D. A. (1940). Plant microtechnique. McGraw-Hill Book Co. New York and London. pp. 126-154.
- Karut, K.; C. Kazak; I. Doker and A. A. Y. Malik (2012). Natural parasitism of *Bemisia tabaci* (Hemiptera: Aleyrodidae) by native Aphelinidae (Hymenoptera) parasitoids in tomato greenhouses in Mersin, Turkey. IOBC/WPRS Bulletin. 80: 69-74.
- Leite, G. L. D.; M. Picanco; A. A. Azevedo and A. H. R. Gonring (1999). Effect of trichomes, allelochemicals and minerals on the resistance of *Lycopersicon hirsutum* to tomato leaf miner. Pesquisa Agropecuaria Brasileira, 34 (11): 2059-2064.
- Leite, G. L. D.; M. Picanco; G. N. Jham and F. Marquini (2004). Intensity of *Tuta absoluta* (Meyrick, 1917) (Lepidoptera: Gelechiidae) and *Liriomyza* spp. (Diptera: Agromyzidae) attacks on *Lycopersicon esculentum* Mill leaves. Ciénciae Agrotecnologia, 28 (1): 42-48.
- Leite, G. L. D.; M. Picanco; G. N. Jham and M. D. Moreira (2004). Natural factors influencing whitefly attack in tomato. Arquivos do Instituto Biológico (Sao Paulo), 71 (2): 245-248.
- Marcos, H.; R. Jordi; D. Rios; R. Elena and D. Carlos (2005). Chemical composition of cultivar of tomatoes resistant and non resistant against the Tomato Yellow Leaf Curl Virus (TYLCV). Electronic J. Environ. Agric. Food Chem., 4 (5): 29-35.
- Millonig, G. (1961). Advantages of a phosphate buffer of  $O_5O_4$  solution in fixation. J. Appl. Phys., 32:1637.
- Mohasin, M.; A. Banerjee; S. Singha and S. Hazra (2005). Influence of abiotic factors and leaf characters on whitefly, *Bemisia tabaci* Genn. (Aleyrodidae: Homoptera), population on different varieties of pumpkin. Horticultural J., 18 (1): 42-45.
- Mukhtar, S. K.; A. Hashim; M. Peterschmitt and M. K. Abdrahman (2009). Field screening and molecular identification of tomato leaf curl virus in Sudan source. Arab J. Plant Protection, 27(1): 95-98.
- Nannini, M.; F. Atzori; F. Foddi; R. Pesci and F. Sanna (2011). A Survey of *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) Outbreaks in Tomato Greenhouses in Southern Sardinia (Italy). Acta Hort. 917:39-46.
- Natarajan, K. (1987). Influence of nitrogen, phosphorus, potassium fertilization on the population density of cotton whitefly, India. Illus. Paper, 134-136.
- Oliveira, A. C. R.; V. R. S. Veloso; R. G. Barros; P. M. Fernandes and E. R. B. Souza (2008). Capture of *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) with light trap in tomato crop. Pesquisa Agropecuaria Tropical, 38 (3): 153-157.
- Rote, N. B. and S. N. Puri (1992). Effects of fertilizer application on incidence of whitefly on different cotton cultivars. J. Maharashtra Agric. Univ., 17: 45-48.

- Sahu, I. K. and S. S. Shaw (2005). Relative susceptibility of different tomato genotypes by whitefly, *Bemisia tabaci* Gen. J. Interacademia, 9 (3): 376-381.
- Samarajeewa, P. K.; M. G. M. K. Meegahakumbura; R. M. S. L. Rajapakse; C. G. Gammulla and V. A. Sumanasinghe (2005). Molecular and morphological identification of Tomato Yellow Leaf Curl Virus (TYLCV) disease in tomato. Ann. SriLanka Dept. Agric., 7: 233-244.
- Setiawati, W; B. K. Udiarto and N. Gunaeni (2009). Preference and infestation pattern of *Bemisia tabci* (Genn.) on some tomato varieties and its effect on gemini virus infestation. Indonesian J. Agric., 2 (1): 57-64.
- Snyder, J. and C. Min (2012). Insect resistance in *Lycopersicon hirsutum* LA2329-current status. Acta Horticulturae, (944): 15-21.
- Spasov, D.; D. Spasova and B. Atanasova (2013). Harmfulness and population dynamics of *Tuta absoluta* (Meyrick,1917) in Strumica region. Zbornik Radova. 18 (20):159-163.
- Summers, C. G.; A. S. Newton; J. P. Mitchell and J. J. Stapleton (2010). Population dynamics of arthropods associated with early-season tomato plants as influenced by soil surface microenvironment. Crop Protection, 29: 249-254.
- Vuong, P. T.; J. Kim and Y. Song (2001). The seasonal occurrence of the two Aphid species, *Myzus persicae* and *Aphis gossypii*, and their natural enemies on vegetable crops in Chinju, Korea. J. Asia-Pacific Entomol., 4 (1): 41-44.
- Wen Feng F.; Z. Peng; T. Jinchang; Z. Yun Ding; B. Shou Dong; G. Ji Guang and X. ZengEn (2009). Temporal and spatial relationships among *Bemisia tabaci* and its natural enemies in tomato fields. J. China Agric. Univ., 14 (4): 77-83.

### تقييم صنفين من الطماطم للإصابة ببعض الآفات الحشرية يوسف عز الدين يوسف عبد الله و هناء صالح أبو بكر فرج قسم وقاية النبات - كلية الزراعة - جامعة عين شمس

أجريت التجارب في حقل طماطم بمحافظة القليوبية - مصر زرع بصنفين من الطماطم لاختبار حساسيتهما للإصابة بحشرات ذبابة الطماطم الأبيض والممن ونطاط أوراق البطاطس وحافرة أوراق الطماطم بالإضافة لدراسة التواجد الموسمي لهذه الآفات على الصنفين خلال العروة الصيفية لموسمي ٢٠١٢ ، ٢٠١٣. تشير النتائج إلى أن الذبابة البيضاء والممن كان لكل منهما قمتين للتواجد خلال مايو ويونيو في كلا الموسمين وعلى كلا الصنفين. بالنسبة لنطاط أوراق البطاطس فكان له أيضا قمتين للتواجد خلال نفس الشهرين ولم يختلف الصنفين معنويا في شدة الإصابة به. كان تواجد الأطوار المختلفة لحشرة ناخرة أوراق الطماطم متساويا تقريبا على كلا الصنفين في موسم ٢٠١٢ بينما لم تتواجد تقريبا خلال موسم ٢٠١٣. كان الارتباط بين شدة الإصابة بكل الحشرات موضع الدراسة وعدد الشعرات الموجودة على أوراق الطماطم مأخوذة من الصنفين في ٢٠ مايو ، ٤ يونيو سالبًا وعالي المعنوية فيما عاد الممن الذي اختفى تماما خلال أشهر الصيف من على الصنف Hybrid Super. في التاريخ الثالث للعينة (٢٠ يونيو، ٢٠١٣) كان الارتباط سالبًا وعالي المعنوية أيضا مع تعداد حشرات الذبابة البيضاء بينما كان موجبا مع تعداد أطوار ناخرة أوراق الطماطم ولوحظت نفس النتائج على الصنف Crystal HYB. كانت تأثيرات عناصر النيتروجين، الفوسفور والكالسيوم موجبة وعالية المعنوية على تعداد كل الآفات موضع الدراسة على كلا الصنفين بينما كان تأثير عنصر البوتاسيوم سالبًا وعالي المعنوية. كان تأثير الماغنسيوم موجبا وشديد المعنوية على الآفات المختلفة على الصنف Hybrid بينما كان الارتباط سالبًا مع الآفات الثاقبة الماصة على الصنف Crystal. تشير النتائج أيضا إلى أن الحديد، الزنك والمنجنيز كان تأثيرهم موجبا وشديد المعنوية بينما كان تأثير النحاس سالبًا على تواجد الآفات موضع الدراسة على الصنفين.